



**Check with local regulatory before use**

# PRODUCT BOOK

Exclusively for veterinarians & veterinary clinics

JUNE 2019



**VETERINARY HEALTH NUTRITION**



# WHO WE ARE

In 1968 in the South of France, a veterinarian named Jean Cathary discovered he could help dogs heal more fully through tailored nutrition. In this visionary act, ROYAL CANIN® was born. Today, using advanced veterinary science and careful observation to identify the most adapted nutrients, our 220+ individual formulas support the health of every individual cat and dog.



# GENERAL INFORMATION

## WHAT SETS US APART

ROYAL CANIN® recognizes that there are fundamental differences between breeds, sizes, and life stages that go well beyond physical appearance. Through science and research we have gained a profound understanding of these differences. We apply our knowledge to developing innovative nutritional solutions that respect each individual pet's needs. At ROYAL CANIN®, we obsess over details, providing tailored nutrition to magnify cats and dogs beauty.

## VETERINARY HEALTH NUTRITION : THE NEW RANGE

Organized around 6 core pet health needs to assist in the nutritional management of pets with the following issues:

### HEALTH MANAGEMENT

Optimal weight  
Senior health  
Oral hygiene  
Periods of adaptation

### URINARY

Lower urinary tract disease

### DERMATOLOGY

Adverse food reaction  
Dermatosis & hair loss

### WEIGHT MANAGEMENT

Overweight / Obesity  
Diabetes Mellitus

### VITAL SUPPORT

Renal disease  
Mobility  
Heart failure

### GASTROINTESTINAL TRACT

Gastrointestinal disorders  
Liver disease  
Convalescence

## EASY TO USE PRODUCT GUIDE

This product guide is an easy reference to the new ROYAL CANIN VETERINARY Health Nutrition range. Each diet listing provides product information, feeding recommendations, ingredient listings, nutrient profiles and package availability. This guide is a valuable resource to help you select the best nutritional solutions for your canine and feline patients.

**Product name**

**URINARY S/O**

**LOWER URINARY TRACT DISEASE**

**RECOMMENDED FOR CASES OF**

- Struvite crystals: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence

**NOT RECOMMENDED IN CASE OF**

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine acidifying drugs
- Growth, gestation/lactation

**STRUVITE DISSOLUTION**  
Helps dissolve all types of struvite stones.

**URINE DILUTION**  
Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.

**LOW PSS**  
Helps lower the concentration of ions contributing to crystal formation.

**LOW MAGNESIUM**  
Reduced level of magnesium, a natural component of struvite crystals.

**KEY VALUES**

	Per 1000 kcal	Per 100g as fed
Protein	81.1 g	24.5 g
Fat content	38.7 g	15 g
Carbohydrate	85.7 g	22.2 g
Cruke fibre	15 g	2.9 g
Dietary fibre	17.5 g	4.5 g
Calcium	2.3 g	0.9 g
Phosphorus	2.3 g	0.9 g
Potassium	2.6 g	1 g
Sodium	3.4 g	1.30 g
Chloride	3.8 g	2.25 g
Magnesium	0.1 g	0.05 g
Copper	5.9 mg	1.5 mg
Zinc	43.9 mg	17 mg
Taurine	0.6 g	0.23 g
Vitamin A	4198.3 IU	2400 IU
Vitamin E	121.7 mg	51 mg
Omega 6	8.7 g	3.36 g
Omega 3	2.3 g	0.88 g
EPA + DHA	1 g	0.37 g
Metabolisable energy*		387 kcal / 100g

**RECOMMENDED DAILY INTAKE & FEEDING DURATION**

CAT'S WEIGHT (kg)	BODY CONDITION					
	THIN		NORMAL		OVERWEIGHT	
	grams	cups	grams	cups	grams	cups
2	31	4/8	33	3/8	36	2/8
2.5	44	4/8	38	4/8	31	3/8
3	52	5/8	44	4/8	35	3/8
3.5	59	5/8	49	5/8	39	4/8
4	64	6/8	54	5/8	43	4/8
4.5	70	6/8	58	5/8	47	4/8
5	75	7/8	63	4/8	50	5/8
5.5	81	1	67	4/8	54	5/8
6	86	1	72	7/8	57	5/8
6.5	91	1	76	7/8	61	4/8
7	94	1 + 1/8	80	7/8	64	4/8
7.5	101	1 + 1/8	84	1	67	4/8
8	105	1 + 2/8	88	1	70	4/8
8.5	110	1 + 2/8	92	1 + 1/8	73	7/8
9	115	1 + 3/8	96	1 + 1/8	76	7/8
9.5	119	1 + 3/8	99	1 + 1/8	79	7/8
10	123	1 + 3/8	103	1 + 2/8	82	1

**ADULT CAT**

Urinary S/O is a complete dietetic food for cats, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties and its low level of magnesium.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. Then the cat should be re-evaluated regularly by the veterinarian to assess the success of the feeding recommendation. Water should be available at all times.

**NOTE:** Dissolution of struvite stones is not recommended for recurrent idiopathic cystitis, renal function should be checked before the advice of a new diet. In older cats the use of Urinary S/O is not recommended.

**COMPOSITION**  
Rice, selected gluten-free, hydrolysed poultry protein, maize flour, animal fats, hydrolysed animal proteins, maize gluten, minerals, vegetable fibres, fish oil, soya oil, fructo-oligo-saccharides, marigold extract (source of lutein).

**ADDITIVES (kg)\*\*\***  
Nutritional additives: Vitamin A: 21000 IU, Vitamin D3: 800 IU, E1 (Iron): 27 mg, E2 (Zinc): 37 mg, E3 (Copper): 11 mg, E5 (Magnesium): 48 mg, E6 (Zinc): 136 mg, E8 (Biotin): 0.06 mg, Urine acidifiers: 100 mg, Preservatives: Antioxidants.

**Indications & contraindications**

**Nutritional benefits**

**Nutritional informations**

**Ingredients & additives**

## GENERAL FEEDING INSTRUCTIONS

- For consistent and accurate food allocation, it is recommended to weigh your pet's daily food
  - Daily feeding recommendations may be divided into two to four meals
  - Fresh water should be available at all times
  - Individual requirements may vary depending on breed, age, gender, environment, activity level and spay/neuter status
- Mix feeding recommendations are communicated when product performance meet the expectations.

Please note that all indications throughout this book are for dietary support of the listed conditions and do not preclude appropriate medical management. The indications and contraindications are not all-inclusive. For dietary consultations please contact our Technical Services Department.



# TABLE OF CONTENTS

## FELINE

### URINARY

MONOFUNCTION	URINARY S/O	Dry / Wet	
	URINARY S/O MODERATE CALORIE	Dry / Wet	

### DERMATOLOGY

MONOFUNCTION	ANALLERGENIC	Dry	
	HYPOALLERGENIC	Dry	
	SENSITIVITY CONTROL	Dry / Wet	
	SKIN & COAT	Dry / Wet	

### WEIGHT MANAGEMENT

MONOFUNCTION	SATIETY	Dry / Wet	
	DIABETIC	Dry / Wet	

## CANINE

### URINARY

MONOFUNCTION	URINARY S/O	Dry / Wet	
	URINARY S/O AGEING 7+	Dry / Wet	
	URINARY S/O MODERATE CALORIE	Dry / Wet	
	URINARY S/O SMALL DOG	Dry	
	URINARY S/O U/C	Dry	

### DERMATOLOGY

MONOFUNCTION	ANALLERGENIC	Dry	
	HYPOALLERGENIC	Dry / Wet 200 g - 400 g	
	HYPOALLERGENIC MODERATE CALORIE	Dry	
	HYPOALLERGENIC SMALL DOGS	Dry	
	SENSITIVITY CONTROL	Dry / Wet (2)	
	SKIN CARE	Dry	
	SKIN CARE SMALL DOGS	Dry	
	SKIN CARE SMALL DOGS PUPPY	Dry	
	SKIN SUPPORT	Dry	

### WEIGHT MANAGEMENT

MONOFUNCTION	SATIETY	Dry / Wet	
	SATIETY SMALL DOGS	Dry	
	DIABETIC	Dry	
	DIABETIC SPECIAL LOW CARBOHYDRATE	Wet	







# FELINE

VETERINARY HEALTH NUTRITION



# INDEX OF CAT HEALTH ISSUES

PLEASE NOTE: All dietary recommendations throughout this book are for the nutritional management of cats exhibiting listed conditions and do not preclude appropriate medical management. All clinical diets should be used under veterinary supervision.

A	Adverse Food Reactions (AFR: Food allergy/Food intolerance)			-ANALLERGENIC feline	
				-HYPOALLERGENIC feline	
				-SENSITIVITY CONTROL feline	
	Ascites	With heart failure		-RENAL feline	
		With hypoproteinemia	Lymphangiectasia (protein losing enteropathy)	- SENSITIVITY CONTROL feline	
				-GASTRO-INTESTINAL MODERATE CALORIE feline	
			Liver disease	-HEPATIC feline	
	Protein losing nephropathy		-RENAL feline		
Atopy			see dermatitis		
C	Cachexia			-CONVALESCENCE SUPPORT INSTANT	
				-RECOVERY feline	
				-GASTRO-INTESTINAL feline	
	Cancer			see neoplasia	
	Chronic Kidney Disease (CKD)			-RENAL feline	
		Concomitant with Adverse Food reactions		-MULTIFUNCTION feline RENAL-HYPOALLERGENIC	
		Early CKD (IRIS stage 1)		-VCN SENIOR CONSULT feline STAGE 2	
		Azotemic CKD (IRIS stages 2 to 4)		-RENAL FELINE	
	Colitis			see diarrhea	
	Constipation	Responsive to dietary fibre		-FIBRE RESPONSE feline	
		Non responsive to dietary fibre		-GASTRO-INTESTINAL feline recommendation: wet	
	Convalescence			-CONVALESCENCE SUPPORT INSTANT	
				-RECOVERY feline	
			-REHYDRATION SUPPORT INSTANT		
			-GASTRO-INTESTINAL feline		
D	Dermatitis	Cutaneous Adverse Food Reactions (CAFR)		-ANALLERGENIC feline	
				-HYPOALLERGENIC feline	
				-SENSITIVITY CONTROL feline	
				-ANALLERGENIC feline	
				-HYPOALLERGENIC feline	
				-MULTIFUNCTION feline RENAL-HYPOALLERGENIC	
			Concomitant with Chronic Kidney Disease		-MULTIFUNCTION feline RENAL-HYPOALLERGENIC
			Concomitant with Struvite or Calcium Oxalate urolithiasis		-MULTIFUNCTION feline URINARY S/O + HYPOALLERGENIC
		Environmental atopy			-SKIN & COAT feline
					-HYPOALLERGENIC feline
					-ANALLERGENIC feline
		Diabetes Mellitus	During stabilization phase (until control of clinical signs and hyperglycemia) and for patients with BCS ≤6/9		-DIABETIC feline
	For stabilized patients with BCS ≥7/9		-SATIETY feline		
	Diarrhea	Acute diarrhea		-GASTRO-INTESTINAL feline	
				-REHYDRATION SUPPORT INSTANT DIET	
		Chronic diarrhea	Non-specific or unknown etiology		-GASTRO-INTESTINAL feline
					-REHYDRATION SUPPORT INSTANT DIET
		Group of chronic enteropathies / inflammatory bowel disease	Food responsive diarrhea	-ANALLERGENIC feline	
				-HYPOALLERGENIC feline	
				-SENSITIVITY CONTROL feline	
			Antibiotic responsive diarrhea: (in adjunction to adequate antibiotherapy)	-GASTRO-INTESTINAL feline	
				Immunosuppressive drug responsive diarrhea : (in adjunction to anti-inflammatories/ immunosuppressors)	-GASTRO-INTESTINAL feline
					-ANALLERGENIC feline
					-HYPOALLERGENIC feline
			-SENSITIVITY CONTROL feline		
		Lymphangiectasia (protein losing enteropathy)		-SENSITIVITY CONTROL feline	
				-GASTRO-INTESTINAL MODERATE CALORIE feline	
		Small intestinal bacterial overgrowth (SIBO)		-GASTRO-INTESTINAL feline	
	Enteritis associated with hepatitis/pancreatitis			see triaditis	

PLEASE NOTE: The following are recommendations only. The veterinary diets should be used under veterinary supervision.

E	Enteropathies (chronic enteropathies)			see diarrhoea (chronic diarrhoea)
	Exocrine pancreatic insufficiency (EPI): (in adjunction to digestive enzyme treatment)			-GASTRO-INTESTINAL feline
				-HYPOALLERGENIC feline
				-ANALLERGENIC feline
F	FIC (Feline Idiopathic Cystitis) (together with environmental measures)			-MULTIFUNCTION feline URINARY S/O + CALM DRY OR WET
				-URINARY S/O wet feline
	Food allergies/ Food intolerances			see Adverse Food Reactions
	Food elimination trial			-HYPOALLERGENIC feline
				-ANALLERGENIC feline
			-SENSITIVITY CONTROL feline	
G	Gastric disorders	Delayed gastric emptying		-GASTRO-INTESTINAL feline MODERATE CALORIE
		Gastritis, gastric and/or duodenal ulcer		-HYPOALLERGENIC feline
				-ANALLERGENIC feline
				-SENSITIVITY CONTROL feline
				-GASTRO-INTESTINAL feline
Gastro-enteritis			see diarrhea	
H	Heart disease/failure			-RENAL feline
	Hepatic disease	Hepatitis	Acute hepatitis	-RECOVERY AND GASTRO-INTESTINAL feline
			Chronic hepatitis	-HEPATIC feline
			If associated with enteritis/pancreatitis	see triaditis
			Portosystemic shunt	-HEPATIC feline
		Hepatic encephalopathy	-HEPATIC feline	
		Hepatic lipidosis		-RECOVERY
				-CONVALESCENCE SUPPORT INSTANT
		If signs of encephalopathy		-HEPATIC feline
	Hyperlipidemia (fasting hyperlipidemia)			-SATIETY feline
				-VCN SENIOR CONSULT STAGE 1 feline Dry
	Hyperthyroidism (in adjunction to antithyroid treat- ment/ Radioactive iodine)	With chronic kidney disease (CKD)		No specific therapeutic diet VCN Senior Consult Stage 2 High Cal
			-RENAL feline	
Hypertension			-RENAL feline	
I	Idiopathic cystitis (FIC: Feline Idiopathic Cystitis)			see Cystitis
K	Kidney disease			see CKD (Chronic Kidney Disease)
L	Liver disease			see Hepatic disease
M	Malabsorption/ Maldigestion			-GASTRO-INTESTINAL feline
				-ANALLERGENIC feline
				-HYPOALLERGENIC feline
	Malnutrition			see Cachexia
	Megacolon			-FIBRE RESPONSE feline (with cautious follow-up)
				-GASTRO-INTESTINAL feline
	Megaoesophagus	Gastrotomy tube		-CONVALESCENCE SUPPORT INSTANT
				-RECOVERY
			-RECOVERY	
			-GASTRO-INTESTINAL wet feline	
N	Neoplasia (optimal nutrition during cancer treatment)			-RECOVERY
				-GASTRO-INTESTINAL feline
O	Obesity/Overweight (weight loss and weight mainte- nance after weight loss)			-SATIETY feline
	Associated with urolithiasis			-MULTIFUNCTION feline S/O-Satiety
	Œsophagitis			-RECOVERY
	Oral hygiene			-DENTAL feline
	Osteo-articular disorders			-MOBILITY feline



PLEASE NOTE: The following are recommendations only. The veterinary diets should be used under veterinary supervision.

P	Pancreatitis	Acute pancreatitis		-GASTRO-INTESTINAL FELINE MODERATE CALORIE
		Chronic pancreatitis		-GASTRO-INTESTINAL FELINE MODERATE CALORIE
			If complicated with diabetes	-DIABETIC feline
		If associated with enteritis/hepatitis		see triaditis
	Poor appetite		-GASTRO-INTESTINAL feline	
			-RECOVERY	
Pre/post-surgery: (including digestive surgery)		-CONVALESCENCE SUPPORT INSTANT		
		-GASTRO-INTESTINAL feline		
		-REHYDRATION SUPPORT		
		-RECOVERY		

R	Renal disease/failure	see CKD (Chronic Kidney Disease)
---	-----------------------	----------------------------------

S	Stress-and anxiety-related disorders	-CALM feline
	When associated with urinary signs	-MULTIFUNCTION feline URINARY S/O + CALM

T	Tartar and dental plaque	-DENTAL feline
---	--------------------------	----------------


	Triaditis (combination of enteritis, cholangitis or cholangiohepatitis, and pancreatitis)	Stable cats/ long-term management	-HYPOALLERGENIC feline	
			-ANALLERGENIC feline	
			-SENSITIVITY CONTROL feline	
			-GASTRO-INTESTINAL feline	
		Cats with dysorexia and/ or needing intensive care or convalescence	-RECOVERY	
		Except in case of hepatic encephalopathy, then	-HEPATIC feline	
	Tube feeding	TO UPDATE with liquid diets: Tube with diameter <10Fr		
		Need of protein and/or phosphorus restriction (kidney disease, hepatic encephalopathy...)	-RENAL LIQUID feline	
			All other conditions	-RECOVERY LIQUID -CONVALESCENCE SUPPORT INSTANT
		Tube with diameter ≥10 Fr	Non-specific conditions	-RECOVERY LIQUID OR WET -CONVALESCENCE SUPPORT INSTANT
			Need of protein and/or phosphorus restriction (kidney disease, hepatic encephalopathy...)	-RENAL LIQUID feline
		Other specific conditions: corresponding specific diet mixed with water and blended (wet diet with same volume of water,or kibbles soaked with water*)		

U	Urolithiasis	Struvite urolithiasis (prevention & dissolution)	-URINARY S/O feline
---	--------------	--	---------------------

	Associated with Adverse Food Reactions	-MULTIFUNCTION feline URINARY S/O + HYPOALLERGENIC	
		Associated with overweight/obesity	-MULTIFUNCTION feline URINARY S/O + SATIETY
		Associated with stress-related disorders	-MULTIFUNCTION feline URINARY S/O + CALM
		Calcium oxalate and calcium phosphate urolithiasis (prevention)	-URINARY S/O feline
	Associated with Adverse Food Reactions	-MULTIFUNCTION feline S/O + HYPOALLERGENIC	
		Associated with overweight/obesity	-MULTIFUNCTION feline URINARY S/O + SATIETY
		Associated with stress	-MULTIFUNCTION feline URINARY S/O + CALM
		Calcium oxalate nephroliths associated with Chronic Kidney Disease (CKD)	-RENAL feline
	Purine (Urate, xanthine) urolithiasis (prevention)		-HYPOALLERGENIC dry feline and try to increase water consumption**
	Cystine urolithiasis (prevention)	-RENAL feline wet loaf or chunks in gravy beef or tuna	
		-HYPOALLERGENIC dry and try to increase water consumption**	

\* Kibbles rehydration for blending  
- mix one volume of kibbles with 3 volumes of hot tap water (around 60°)  
- let the kibbles absorb water for about 15 min before blending

\*\* Increasing water consumption in cats (cats being less prone than dogs to consume rehydrated kibbles) Water-related factors to consider include freshness, taste, movement (water fountains, dripping faucets, bowl with pump-bubbled air...), and shape of the container.





## URINARY S/O



Available in 400 g, 1.5 kg, 3.5 kg, 7 kg, 9 kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

CAT'S WEIGHT (kg)	BODY CONDITION					
	THIN		NORMAL		OVERWEIGHT	
	grams	cups	grams	cups	grams	cups
2	39	4/8	33	3/8	26	2/8
2,5	46	4/8	38	4/8	31	3/8
3	52	5/8	44	4/8	35	3/8
3,5	59	5/8	49	5/8	39	4/8
4	64	6/8	54	5/8	43	4/8
4,5	70	6/8	58	5/8	47	4/8
5	75	7/8	63	6/8	50	5/8
5,5	81	1	67	6/8	54	5/8
6	86	1	72	7/8	57	5/8
6,5	91	1	76	7/8	61	6/8
7	96	1 + 1/8	80	7/8	64	6/8
7,5	101	1 + 1/8	84	1	67	6/8
8	105	1 + 2/8	88	1	70	6/8
8,5	110	1 + 2/8	92	1 + 1/8	73	7/8
9	115	1 + 3/8	96	1 + 1/8	76	7/8
9,5	119	1 + 3/8	99	1 + 1/8	79	7/8
10	123	1 + 3/8	103	1 + 2/8	82	1

Urinary S/O is a complete dietetic feed for cats, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties and its low level of magnesium.

RECOMMENDATIONS: It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. Then the cat should be re-evaluated regularly by the veterinarian to assess the relevance of a life long recommendation. Water should be available at all times.

**NOTE:** - the wet version of Urinary S/O Feline is recommended for recurrent idiopathic cystitis,  
- renal function should be checked before the advice of a new diet. In older cats the use of Urinary S/O Ageing<sup>®</sup> + Calm is recommended.

## LOWER URINARY TRACT DISEASE

### RECOMMENDED FOR CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence

### NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying drugs
- Growth, gestation/lactation



#### STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



#### URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



#### LOW RSS

Helps lower the concentration of ions contributing to crystal formation.



#### LOW MAGNESIUM

Reduced level of magnesium, a natural component of struvite crystals.

**[S/O®] INDEX** This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	89.1 g	34.5 g
Fat content	38.7 g	15 g
Carbohydrate	85.7 g	33.2 g
Crude fibre	7.5 g	2.9 g
Dietary fibre	17.6 g	6.8 g
Calcium	2.3 g	0.9 g
Phosphorus	2.3 g	0.9 g
Potassium	2.6 g	1 g
Sodium	3.4 g	1.30 g
Chloride	5.8 g	2.26 g
Magnesium	0.1 g	0.05 g
Copper	3.9 mg	1.5 mg
Zinc	43.9 mg	17 mg
Taurine	0.6 g	0.23 g
Vitamin A	6198.3 IU	2400 IU
Vitamin E	131.7 mg	51 mg
Omega 6	8.7 g	3.38 g
Omega 3	2.3 g	0.88 g
EPA + DHA	1 g	0.37 g
Metabolisable energy*		387 kcal / 100 g

\*Calculated according to NRC2006.

### COMPOSITION

Rice, wheat gluten\*\*, dehydrated poultry protein, maize flour, animal fats, hydrolysed animal proteins, maize gluten, minerals, vegetable fibres, fish oil, soya oil, fructo-oligo-saccharides, marigold extract (source of lutein).

\*\*L.I.P.: protein selected for its very high digestibility

### ADDITIVES (KG)\*\*\*

Nutritional additives: Vitamin A: 21500 IU, Vitamin D3: 800 IU, E1 (Iron): 37 mg, E2 (Iodine): 3.7 mg, E4 (Copper): 11 mg, E5 (Manganese): 48 mg, E6 (Zinc): 136 mg, E8 (Selenium): 0.06 mg, Urine acidifying substances: calcium sulphate (1.25%). Preservatives - Antioxidants.

\*\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



## URINARY S/O

WEIGHT MANAGEMENT

GASTROINTESTINAL TRACT

VITAL SUPPORT

HEALTH MANAGEMENT

CANINE



## URINARY S/O



Available in box of 12 pouches of 85 g

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT CAT	CAT'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	pouch	grams	pouch	grams	pouch
	2	170	2	140	1 +1/2	110	1 +1/2
	2,5	195	2 +1/2	165	2	130	1 +1/2
	3	225	2 +1/2	185	2	150	2
	3,5	250	3	210	2 +1/2	165	2
	4	275	3	230	2 +1/2	185	2
	4,5	300	3 +1/2	250	3	200	2 +1/2
	5	325	4	270	3	215	2 +1/2
	5,5	345	4	290	3 +1/2	230	2 +1/2
	6	365	4 +1/2	305	3 +1/2	245	3
	6,5	390	4 +1/2	325	4	260	3
	7	410	5	340	4	275	3
	7,5	430	5	360	4	285	3 +1/2
	8	450	5 +1/2	375	4 +1/2	300	3 +1/2
	8,5	470	5 +1/2	390	4 +1/2	315	3 +1/2
	9	490	6	410	5	325	4
	9,5	510	6	425	5	340	4
	10	530	6	440	5	350	4

Urinary S/O is a complete dietetic feed for cats, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties and its low level of magnesium.

RECOMMENDATIONS: It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. Then the cat should be re-evaluated regularly by the veterinarian to assess the relevance of a life long recommendation. Water should be available at all times.

NOTE: renal function should be checked before the advise of a new diet. In older cats the use of urinary S/O Ageing<sup>®</sup> + Calm is recommended.

## LOWER URINARY TRACT DISEASE

### RECOMMENDED FOR CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Feline idiopathic cystitis (FIC)

### NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying drugs
- Growth, gestation/lactation



#### IDIOPATHIC CYSTITIS

Thanks to the high moisture content of wet food, Urinary S/O Feline increases urine dilution. Since nearly 64% of cases of FLUTD\* are idiopathic cystitis, the wet diet should be preferred in cases of recurrent cystitis.



#### STRUVITE DISSOLUTION

Urinary S/O Feline effectively dissolves struvite uroliths.



#### LOW RSS

Undersaturated urine is an environment where crystals cannot proliferate. Thus the urine is unfavourable to the development of struvite and calcium oxalate crystals.



#### URINE DILUTION

Diluting urine decreases the urinary concentration of struvites and calcium oxalates. The larger urinary volume also ensures regular bladder wash-out.

[S/O<sup>®</sup> INDEX] This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	130.18 g	11 g
Fat content	41.42 g	3.5 g
Carbohydrate	33.14 g	2.8 g
Crude fibre	11.83 g	1 g
Dietary fibre	17.75 g	1.5 g
Calcium	2.96 g	0.25 g
Phosphorus	2.96 g	0.25 g
Potassium	2.6 g	0.22 g
Sodium	3.55 g	0.3 g
Chloride	4.73 g	0.4 g
Magnesium	0.19 g	0.02 g
Copper	4.62 mg	0.39 mg
Zinc	30.77 mg	2.6 mg
Taurine	1.18 mg	0.1 g
Vitamin A	26035.5 IU	2200 IU
Vitamin E	118.34 mg	10 mg
Omega 6	8.88 g	0.75 g
Omega 3	1.89 g	0.16 g
EPA + DHA	1.07 g	0.09 g
Metabolisable energy*		85 kcal / 100 g

\*Calculated according to NRC2006.

### COMPOSITION

Meat and animal derivatives, derivatives of vegetable origin, minerals, cereals, vegetable protein extracts, oils and fats, various sugars.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 50 IU, E1 (Iron): 3 mg, E2 (Iodine): 0.3 mg, E4 (Copper): 2.5 mg, E5 (Manganese): 0.9 mg, E6 (Zinc): 9 mg, Urine acidifying substances: calcium sulfate: 0.35%.

Technological additives: (acidity regulator) Sodium bisulfate: 0.5 g.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



## URINARY S/O



## URINARY S/O



Available in box of 12 pouches of 85 g

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT CAT	CAT'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	pouch	grams	pouch	grams	pouch
	2	170	2	140	1 +1/2	110	1 +1/2
	2,5	195	2 +1/2	165	2	130	1 +1/2
	3	225	2 +1/2	185	2	150	2
	3,5	250	3	210	2 +1/2	165	2
	4	275	3	230	2 +1/2	185	2
	4,5	300	3 +1/2	250	3	200	2 +1/2
	5	320	4	270	3	215	2 +1/2
	5,5	345	4	285	3 +1/2	230	2 +1/2
	6	365	4 +1/2	305	3 +1/2	245	3
	6,5	390	4 +1/2	325	4	260	3
	7	410	5	340	4	275	3
	7,5	430	5	360	4	285	3 +1/2
	8	450	5 +1/2	375	4 +1/2	300	3 +1/2
	8,5	470	5 +1/2	390	4 +1/2	315	3 +1/2
	9	490	6	410	5	325	4
	9,5	510	6	425	5	340	4
	10	525	6	440	5	350	4

Urinary S/O is a complete dietetic feed for cats, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties.

RECOMMENDATIONS: It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. Then the cat should be re-evaluated regularly by the veterinarian to assess the relevance of a life long recommendation. Water should be available at all times.

NOTE: renal function should be checked before the advise of a new diet. In older cats the use of Urinary S/O Ageing<sup>®</sup> + Calm is recommended.

## LOWER URINARY TRACT DISEASE

### RECOMMENDED FOR CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Feline idiopathic cystitis (FIC)

### NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying drugs
- Growth, gestation/lactation



#### IDIOPATHIC CYSTITIS

Thanks to the high moisture content of wet food, Urinary S/O Feline increases urine dilution. Since nearly 64% of cases of FLUTD\* are idiopathic cystitis, the wet diet should be preferred in cases of recurrent cystitis.



#### STRUVITE DISSOLUTION

Urinary S/O Feline effectively dissolves struvite uroliths.



#### LOW RSS

Undersaturated urine is an environment where crystals cannot proliferate. Thus the urine is unfavourable to the development of struvite and calcium oxalate crystals.



#### URINE DILUTION

Diluting urine decreases the urinary concentration of struvites and calcium oxalates. The larger urinary volume also ensures regular bladder wash-out.

[S/O<sup>®</sup> INDEX] This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	119.76 g	10 g
Fat content	41.92 g	3.5 g
Carbohydrate	44.31 g	3.7 g
Crude fibre	11.98 g	1 g
Dietary fibre	14.37 g	1.2 g
Calcium	2.63 g	0.22 g
Phosphorus	2.28 g	0.19 g
Potassium	2.16 g	0.18 g
Sodium	3.59 g	0.3 g
Chloride	5.03 g	0.42 g
Magnesium	0.14 g	0.01 g
Copper	4.55 mg	0.38 mg
Zinc	25.15 mg	2.10 mg
Taurine	1.2 g	0.1 g
Vitamin A	35928.14 IU	3000 IU
Vitamin E	147.31 mg	12.30 mg
Omega 6	8.38 g	0.7 g
Omega 3	1.8 g	0.15 g
EPA + DHA	1.1 g	0.09 g
Metabolisable energy*		91 kcal / 100 g

\*Calculated according to NRC2006.

### COMPOSITION

Meat and animal derivatives, cereals, vegetable protein extracts, derivatives of vegetable origin, minerals, oils and fats, various sugars.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 225 IU, E1 (Iron): 3 mg, E2 (Iodine): 0.25 mg, E4 (Copper): 2 mg, E5 (Manganese): 1 mg, E6 (Zinc): 10 mg, Urine acidifying substances: calcium sulfate: 0.25%.

Technological additives: (acidity regulator) Sodium bisulfate: 0.5 g.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



## URINARY S/O





## URINARY S/O

MODERATE CALORIE



Available in 400 g, 1.5 kg, 3.5 kg, 7 kg, 9 kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT CAT	CAT'S WEIGHT (kg)	BODY CONDITION			
		NORMAL		OVERWEIGHT	
		grams	cups	grams	cups
	2	36	3/8	29	3/8
	2,5	42	4/8	34	3/8
	3	48	5/8	39	4/8
	3,5	54	5/8	43	4/8
	4	59	6/8	47	4/8
	4,5	64	6/8	51	5/8
	5	69	7/8	55	5/8
	5,5	74	7/8	59	6/8
	6	79	1	63	6/8
	6,5	84	1	67	6/8
	7	88	1	70	7/8
	7,5	92	1 + 1/8	74	7/8
	8	97	1 + 1/8	77	7/8
	8,5	101	1 + 2/8	81	1
	9	105	1 + 2/8	84	1
	9,5	109	1 + 2/8	87	1
	10	113	1 + 3/8	91	1 + 1/8

Urinary S/O Moderate Calorie is a complete dietetic feed for cats, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties and its low level of magnesium.

RECOMMENDATIONS : It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O Moderate Calorie for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. Then the cat should be re-evaluated regularly by the veterinarian to assess the relevance of a life long recommendation. Water should be available at all times.

NOTE: in older cats, renal function should be checked before Urinary S/O Moderate Calorie is advised.  
The wet version of Urinary S/O Moderate Calorie is recommended for recurrent idiopathic cystitis.

## LOWER URINARY TRACT DISEASE

### RECOMMENDED FOR CATS WITH TENDENCY TO BE OVERWEIGHT IN CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence

### NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying drugs
- Growth, gestation/lactation



#### STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



#### MODERATE CALORIE

Moderate calorie content to help maintain ideal weight.



#### URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



#### LOW RSS

Helps lower the concentration of ions contributing to crystal formation.

[S/O®] This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	96.73 g	34 g
Fat content	31.29 g	11 g
Carbohydrate	95.87 g	33.7 g
Crude fibre	19.91 g	7 g
Dietary fibre	36.98 g	13 g
Calcium	2.84 g	1 g
Phosphorus	2.56 g	0.9 g
Potassium	2.84 g	1 g
Sodium	3.41 g	1.2 g
Chloride	6.09 g	2.14 g
Magnesium	0.17 g	0.06 g
Copper	4.27 mg	1.5 mg
Zinc	48.65 mg	17.1 mg
Taurine	0.63 g	0.22 g
Vitamin A	6543.39 IU	2300 IU
Vitamin E	119.49 mg	42 mg
Omega 6	7.28 g	2.56 g
Omega 3	1.99 g	0.7 g
EPA + DHA	0.91 g	0.32 g
Metabolisable energy*		352 kcal / 100 g

\*Calculated according to NRC2006.

### COMPOSITION

Rice, dehydrated poultry protein, wheat gluten\*\*, maize flour, vegetable fibres, maize gluten, hydrolysed animal proteins, minerals, animal fats, fish oil, soya oil, fructo-oligo-saccharides, marigold extract (source of lutein).

\*\*L.I.P.: protein selected for its very high digestibility

### ADDITIVES (KG)\*\*\*

Nutritional additives: Vitamin A: 21500 IU, Vitamin D3: 800 IU, E1 (Iron): 35 mg, E2 (Iodine): 3.5 mg, E4 (Copper): 11 mg, E5 (Manganese): 46 mg, E6 (Zinc): 134 mg, E8 (Selenium): 0.06 mg, Urine acidifying substances: calcium sulphate (1.3%). Preservatives - Antioxidants.

\*\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



URINARY S/O MODERATE CALORIE



## URINARY S/O

MODERATE CALORIE



MORSELS IN GRAVY

Available in box of 12 pouches of 85 g

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT CAT	CAT'S WEIGHT (kg)	BODY CONDITION			
		NORMAL		OVERWEIGHT	
		grams	pouch	grams	pouch
	2	160	2	130	1 + 1/2
	2,5	185	2	150	2
	3	215	2 + 1/2	170	2
	3,5	240	3	190	2
	4	260	3	210	2 + 1/2
	4,5	285	3 + 1/2	225	2 + 1/2
	5	305	3 + 1/2	245	3
	5,5	330	4	260	3
	6	350	4	280	3 + 1/2
	6,5	370	4 + 1/2	295	3 + 1/2
	7	390	4 + 1/2	310	3 + 1/2
	7,5	410	5	325	4
	8	430	5	340	4
	8,5	445	5	355	4
	9	465	5 + 1/2	370	4 + 1/2
	9,5	485	5 + 1/2	385	4 + 1/2
	10	500	6	400	4 + 1/2

Urinary S/O Moderate Calorie is a complete dietetic feed for cats, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties and its low level of magnesium.

RECOMMENDATIONS: It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O Moderate Calorie for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. Then the cat should be re-evaluated regularly by the veterinarian to assess the relevance of a life long recommendation. Water should be available at all times.

NOTE: in older cats, renal function should be checked before Urinary S/O Moderate Calorie Feline is advised.

## LOWER URINARY TRACT DISEASE

### RECOMMENDED FOR CATS WITH TENDENCY TO BE OVERWEIGHT IN CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Feline idiopathic cystitis (FIC)

### NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying drugs
- Growth, gestation/lactation



#### URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate uroliths.



#### STRUVITE DISSOLUTION

Helps dissolve struvite uroliths.



#### LOW RSS

Helps lower the concentration of ions contributing to crystal formation.



#### MODERATE CALORIE

Moderate calorie content to help maintain ideal weight.

[S/O®] This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	129.96 g	9.5 g
Fat content	36.94 g	2.7 g
Carbohydrate	45.14 g	3.3 g
Crude fibre	17.78 g	1.3 g
Dietary fibre	20.52 g	1.5 g
Calcium	3.01 g	0.22 g
Phosphorus	2.6 g	0.19 g
Potassium	2.19 g	0.16 g
Sodium	3.56 g	0.26 g
Chloride	4.51 g	0.33 g
Magnesium	0.14 g	0.01 g
Copper	4.92 mg	0.36 mg
Zinc	24.62 mg	1.80 mg
Taurine	1.16 g	0.09 g
Vitamin A	68399.45 IU	5000 IU
Vitamin E	136 mg	10 mg
Omega 6	5.06 g	0.37 g
Omega 3	2.33 g	0.17 g
EPA + DHA	1.09 g	0.08 g
Metabolisable energy*		73 kcal / 100 g

\*Calculated according to NRC2006.

### COMPOSITION

Meat and animal derivatives, fish and fish derivatives, cereals, derivatives of vegetable origin, vegetable protein extracts, minerals, oils and fats, various sugars.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 170 IU, E1 (Iron): 2 mg, E2 (Iodine): 0.25 mg, E4 (Copper): 2 mg, E5 (Manganese): 0.5 mg, E6 (Zinc): 5 mg, Urine acidifying substances: calcium sulfate: 0.25%.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



URINARY S/O MODERATE CALORIE



# CATS MIX FEEDING GUIDELINES



**24H MIX: URINARY S/O DRY + URINARY S/O LOAF WET**

CAT'S WEIGHT (kg)	POUCH	WEIGHT CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)
2	1+	21	2/8 cup	14	1/8 cup	6	1/8 cup
3	1+	35	3/8 cup	26	2/8 cup	15	1/8 cup
4	1+	48	4/8 cup	37	3/8 cup	23	2/8 cup
5	1+	61	6/8 cup	47	4/8 cup	30	3/8 cup
6	1+	72	7/8 cup	57	5/8 cup	37	3/8 cup
7	1+	84	1 cup	66	6/8 cup	44	4/8 cup
8	1+	94	1 cup + 1/8 cup	75	7/8 cup	50	5/8 cup
9	1+	105	1 cup + 2/8 cup	84	1 cup	57	5/8 cup
10	1+	115	1 cup + 3/8 cup	93	1 cup + 1/8 cup	63	6/8 cup




**24H MIX: URINARY S/O DRY + URINARY S/O CIG WET**

CAT'S WEIGHT (kg)	POUCH	WEIGHT CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)
2	1+	21	2/8 cup	14	1/8 cup	6	1/8 cup
3	1+	35	3/8 cup	26	2/8 cup	15	1/8 cup
4	1+	48	4/8 cup	37	3/8 cup	23	2/8 cup
5	1+	60	6/8 cup	47	4/8 cup	30	3/8 cup
6	1+	72	7/8 cup	57	5/8 cup	37	3/8 cup
7	1+	84	1 cup	66	6/8 cup	44	4/8 cup
8	1+	94	1 cup + 1/8 cup	75	7/8 cup	50	5/8 cup
9	1+	105	1 cup + 2/8 cup	84	1 cup	57	5/8 cup
10	1+	115	1 cup + 3/8 cup	93	1 cup + 1/8 cup	63	6/8 cup



**24H MIX: URINARY S/O MODERATE CALORIE DRY + URINARY S/O MODERATE CALORIE WET**

CAT'S WEIGHT (kg)	POUCH	WEIGHT CONDITION			
		NORMAL		OVERWEIGHT	
		DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)
2	1+	18	2/8 cup	10	1/8 cup
3	1+	31	3/8 cup	19	2/8 cup
4	1+	43	4/8 cup	28	3/8 cup
5	1+	55	5/8 cup	36	3/8 cup
6	1+	65	6/8 cup	44	4/8 cup
7	1+	76	7/8 cup	51	5/8 cup
8	1+	86	1 cup	58	6/8 cup
9	1+	95	1 cup + 1/8 cup	65	6/8 cup
10	1+	105	1 cup + 2/8 cup	72	7/8 cup



## ANALLERGENIC



Available in 2 kg, 4 kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

	CAT'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	cups	grams	cups	grams	cups
ADULT CAT	2	39	3/8	32	3/8	26	2/8
	2.5	46	4/8	38	3/8	30	3/8
	3	52	4/8	43	4/8	35	3/8
	3.5	58	5/8	48	4/8	39	3/8
	4	64	5/8	53	5/8	42	4/8
	4.5	69	6/8	58	5/8	46	4/8
	5	75	6/8	62	5/8	50	4/8
	5.5	80	7/8	67	6/8	53	5/8
	6	85	7/8	71	6/8	57	5/8
	6.5	90	1	75	6/8	60	5/8
	7	95	1	79	7/8	63	5/8
	7.5	100	1 + 1/8	83	7/8	66	6/8
	8	104	1 + 1/8	87	7/8	69	6/8
	8.5	109	1 + 1/8	91	1	73	6/8
	9	113	1 + 2/8	95	1	76	6/8
	9.5	118	1 + 2/8	98	1	79	7/8
	10	122	1 + 2/8	102	1 + 1/8	81	7/8

When a food allergy or a food intolerance is suspected, the cat's diet should be changed following classical food transition. Nutritional management with Anallergenic may be continued for the pet's lifetime.

## ADVERSE REACTION TO FOOD



### RECOMMENDED FOR CASES OF

- Adverse Food Reactions (AFR) with dermatologic and/or gastro-intestinal signs:
  - diagnosis: food elimination trial
  - management
- Feline Atopic Dermatitis (CAD) associated with AFR
- Inflammatory Bowel Disease (IBD)

### NOT RECOMMENDED IN CASE OF

- Growth, gestation and lactation



#### OLIGOPEPTIDES

Low molecular weight peptides to reduce the risk of adverse food reaction.



#### ALLERGEN RESTRICTION

Formula and production process aimed at excluding sources of food allergens.



#### SKIN BARRIER

Formulated to support the skin's natural protective barrier for optimal skin health.



#### ANTIOXIDANT COMPLEX

A synergistic complex of antioxidants to help neutralise free radicals.



This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	62.9 g	24.6 g
Fat content	43.4 g	17 g
Carbohydrate	106.3 g	41.6 g
Crude fibre	9.5 g	3.7 g
Dietary fibre	20.4 g	8 g
Calcium	1.5 g	0.6 g
Phosphorus	1.2 g	0.48 g
Potassium	1.5 g	0.6 g
Sodium	2.8 g	1.1 g
Chloride	4 g	1.58 g
Magnesium	0.1 g	0.04 g
Copper	3.8 mg	1.5 mg
Zinc	46.8 mg	18.3 mg
Taurine	0.7 g	0.28 g
Vitamin A	8924.2 IU	3650 IU
Vitamin E	150.8 mg	59 mg
Omega 6	9.3 g	3.64 g
Omega 3	2.7 g	1.07 g
EPA + DHA	1.2 g	0.47 g
Metabolisable energy*		409 kcal / 100 g

\*Calculated according to NRC2006.

### COMPOSITION

Maize starch, feather hydrolysate with low molecular weight (source of L amino acids and oligopeptides), copra oil, soya oil, vegetable fibres, minerals, animal fat, fish oil, chicory pulp, fructo-oligo-saccharides, maltodextrin, mono - and diglycerides of palmitic and stearic acids esterified with citric acid, dextrose, marigold extract (source of lutein). Protein source: feather hydrolysate with low molecular weight. Carbohydrate source: maize starch.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin A: 36500 IU, Vitamin D3: 800 IU, E1 (Iron): 42 mg, E2 (Iodine): 3.8 mg, E4 (Copper): 15 mg, E5 (Manganese): 55 mg, E6 (Zinc): 162 mg, E8 (Selenium): 0.26 mg - Technological additives: Clinoptilolite of sedimentary origin: 10 g - Preservatives - Antioxidants.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



## ANALLERGENIC



## HYPOALLERGENIC



Available in 400 g, 2.5 kg, 4.5 kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

	CAT'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	cups	grams	cups	grams	cups
ADULT CAT	2	37	3/8	31	2/8	25	2/8
	2.5	43	3/8	36	3/8	29	2/8
	3	50	4/8	41	3/8	33	3/8
	3.5	55	4/8	46	4/8	37	3/8
	4	61	5/8	51	4/8	41	3/8
	4.5	66	5/8	55	4/8	44	3/8
	5	71	6/8	59	5/8	47	4/8
	5.5	76	6/8	64	5/8	51	4/8
	6	81	6/8	68	5/8	54	4/8
	6.5	86	7/8	72	6/8	57	4/8
	7	90	7/8	75	6/8	60	5/8
	7.5	95	7/8	79	6/8	63	5/8
	8	99	1	83	6/8	66	5/8
	8.5	104	1	87	7/8	69	5/8
	9	108	1	90	7/8	72	6/8
	9.5	112	1 + 1/8	94	7/8	75	6/8
	10	117	1 + 1/8	97	1	78	6/8

When a food allergy or a food intolerance is suspected, the cat's diet should be changed without any food transition. Nutritional management with Hypoallergenic may continue for the cat's lifetime.

## ADVERSE REACTION TO FOOD



### RECOMMENDED FOR CASES OF

- Adverse Food Reactions (AFR) with dermatologic and/or gastro-intestinal signs:
  - diagnosis: food elimination trial
  - management
- Atopic dermatitis associated with AFR
- Chronic diarrhea
- Inflammatory Bowel Disease (IBD)
- Exocrine Pancreatic Insufficiency (EPI)
- Small intestinal bacterial overgrowth (SIBO)

### NOT RECOMMENDED IN CASE OF

- Growth, gestation/lactation



#### HYDROLYSED PROTEIN

Hydrolysed protein with low molecular weight to ensure the food is hypoallergenic.



#### SKIN BARRIER

Formulated to support the skin's natural protective barrier for optimal skin health.



#### EPA/DHA

Fatty acids to help maintain a healthy digestive system and a healthy skin.



#### DIGESTIVE HEALTH

Nutrients which support a balanced digestive system.



This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	62.2 g	25.5 g
Fat content	48.8 g	20 g
Carbohydrate	95.3 g	39.1 g
Crude fibre	8.8 g	3.6 g
Dietary fibre	20 g	8.2 g
Calcium	1.5 g	0.6 g
Phosphorus	1.5 g	0.6 g
Potassium	2 g	0.8 g
Sodium	1.5 g	0.6 g
Chloride	1.9 g	0.79 g
Magnesium	0.1 g	0.05 g
Copper	3.7 mg	1.5 mg
Zinc	41.7 mg	17.1 mg
Taurine	2.5 g	0.25 g
Vitamin A	6582.2 IU	2700 IU
Vitamin E	148.7 mg	61 mg
Omega 6	11.7 g	4.78 g
Omega 3	2.4 g	0.98 g
EPA + DHA	0.8 g	0.32 g
Metabolisable energy*		410 kcal / 100 g

\*Calculated according to NRC2006.

### COMPOSITION

Rice, hydrolysed soya protein isolate, animal fats, vegetable fibres, minerals, beet pulp, hydrolysed poultry liver, soya oil, fish oil, fructo-oligo-saccharides, borage oil, marigold extract (source of lutein). Protein sources: hydrolysed soya protein isolate, hydrolysed poultry liver. Carbohydrate source: rice.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin A: 27000 IU, Vitamin D3: 800 IU, E1 (Iron): 41 mg, E2 (Iodine): 3.7 mg, E4 (Copper): 15 mg, E5 (Manganese): 54 mg, E6 (Zinc): 153 mg, E8 (Selenium): 0.26 mg - Technological additives: Clinoptilolite of sedimentary origin: 5 g - Preservatives - Antioxidants.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



## HYPOALLERGENIC



WEIGHT MANAGEMENT

GASTROINTESTINAL TRACT

VITAL SUPPORT

HEALTH MANAGEMENT

CANINE



## SENSITIVITY CONTROL



Available in 400 g, 1.5 kg, 3.5 kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT CAT	CAT'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	cups	grams	cups	grams	cups
	2	44	4/8	36	3/8	29	2/8
	2.5	51	4/8	43	4/8	34	3/8
	3	58	5/8	49	4/8	39	3/8
	3.5	65	5/8	54	5/8	43	4/8
	4	71	6/8	60	5/8	48	4/8
	4.5	78	7/8	65	5/8	52	4/8
	5	84	7/8	70	6/8	56	5/8
	5.5	90	1	75	6/8	60	5/8
	6	95	1	80	7/8	64	5/8
	6.5	101	1 + 1/8	84	7/8	67	6/8
	7	106	1 + 1/8	89	1	71	6/8
	7.5	112	1 + 1/8	93	1	74	6/8
	8	117	1 + 2/8	98	1	78	7/8
	8.5	122	1 + 2/8	102	1 + 1/8	81	7/8
	9	127	1 + 3/8	106	1 + 1/8	85	7/8
	9.5	132	1 + 3/8	110	1 + 1/8	88	7/8
	10	137	1 + 4/8	114	1 + 2/8	91	1

When a food allergy or a food intolerance is confirmed, the cat's diet should be changed. Nutritional management with Sensitivity Control may continue for the cat's lifetime. Sensitivity Control Feline could be used without any food transition.

## ADVERSE REACTION TO FOOD



### RECOMMENDED FOR CASES OF

- Adverse Food Reactions (AFR) with dermatologic and/or gastrointestinal signs:
  - diagnosis: food elimination trial
  - management
- Chronic diarrhea

### NOT RECOMMENDED IN CASE OF

- Growth, gestation/lactation



#### SELECTED PROTEIN

A limited number of protein sources helps reduce the risk of food allergies.



#### SKIN BARRIER

Formulated to support the skin's natural protective barrier for optimal skin health.



#### EPA/DHA

Fatty acids to help maintain a healthy digestive system and a healthy skin.



#### DIGESTIVE SECURITY

Nutrients which support a balanced digestive system.



**[S/O] INDEX** This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	77.4 g	27 g
Fat content	31.5 g	11 g
Carbohydrate	116.9 g	40.8 g
Crude fibre	17.5 g	6.1 g
Dietary fibre	28.9 g	10.1 g
Calcium	4.6 g	1.61 g
Phosphorus	3.2 g	1.1 g
Potassium	2.3 g	0.8 g
Sodium	2.3 g	0.8 g
Chloride	4 g	1.38 g
Magnesium	0.2 g	0.07 g
Copper	4.3 mg	1.5 mg
Zinc	52.5 mg	18.3 mg
Taurine	0.7 g	0.23 g
Vitamin A	8598.5 IU	3000 IU
Vitamin E	169.1 mg	59 mg
Omega 6	7 g	2.45 g
Omega 3	2.3 g	0.79 g
EPA + DHA	1 g	0.35 g
Metabolisable energy*		349 kcal / 100 g

\*Calculated according to NRC2006.

### COMPOSITION

Rice, dehydrated duck protein, vegetable fibres, hydrolysed poultry proteins, minerals, animal fats, fish oil, soya oil, mono - and diglycerides of palmitic and stearic acids esterified with citric acid, marigold extract [source of lutein]. Protein sources: dehydrated duck protein, hydrolysed poultry proteins. Carbohydrate source: rice.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin A: 26500 IU, Vitamin D3: 800 IU, E1 (Iron): 41 mg, E2 (Iodine): 3.6 mg, E4 (Copper): 15 mg, E5 (Manganese): 53 mg, E6 (Zinc): 150 mg, E8 (Selenium): 0.25 mg - Preservatives - Antioxidants.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

## SENSITIVITY CONTROL



## SENSITIVITY CONTROL

### CHICKEN WITH RICE



Available in box of 12 pouches of 85 g pouch

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT CAT	CAT'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	pouch	grams	pouch	grams	pouch
	2	145	1 + 1/2	125	1 + 1/2	100	1
	2.5	175	2	145	1 + 1/2	115	1 + 1/2
	3	195	2 + 1/2	165	2	130	1 + 1/2
	3.5	220	2 + 1/2	185	2	145	1 + 1/2
	4	240	3	200	2 + 1/2	160	2
	4.5	260	3	220	2 + 1/2	175	2
	5	285	3 + 1/2	235	3	190	2
	5.5	305	3 + 1/2	250	3	200	2 + 1/2
	6	320	4	270	3	215	2 + 1/2
	6.5	340	4	285	3 + 1/2	225	2 + 1/2
	7	360	4	300	3 + 1/2	240	3
	7.5	375	4 + 1/2	315	3 + 1/2	250	3
	8	395	4 + 1/2	330	4	265	3
	8.5	410	5	345	4	275	3
	9	430	5	360	4	285	3 + 1/2
	9.5	445	5	370	4 + 1/2	300	3 + 1/2
	10	465	5 + 1/2	385	4 + 1/2	310	3 + 1/2

Sensitivity Control Feline may be used in case of chronic pruritic dermatosis with an allergic component such as miliary dermatitis.

## ADVERSE REACTION TO FOOD



### RECOMMENDED FOR CASES OF

- Adverse Food Reactions (AFR) with dermatologic and/or gastrointestinal signs:
  - diagnosis: food elimination trial
  - management
- Chronic diarrhea

### NOT RECOMMENDED IN CASE OF

- Growth, gestation/lactation



#### SELECTED PROTEIN

A limited number of protein sources may help reduce the risk of food allergies.



#### SKIN BARRIER

Formulated to support the skin's natural protective barrier for optimal skin health.



#### EPA/DHA

Fatty acids help maintain a healthy digestive system and a healthy skin.



#### DIGESTIVE HEALTH

Adapted nutrients support a balanced digestive system.



**[S/O] INDEX** This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	73.4 g	7.1 g
Fat content	63.1 g	6.1 g
Carbohydrate	50.7 g	4.9 g
Crude fibre	9.3 g	0.9 g
Dietary fibre	14.5 g	1.4 g
Calcium	2.1 g	0.2 g
Phosphorus	2 g	0.19 g
Potassium	2 g	0.19 g
Sodium	1.6 g	0.15 g
Chloride	2.1 g	0.2 g
Magnesium	0.2 g	0.02 g
Copper	4.1 mg	0.4 mg
Zinc	32.1 mg	3.1 mg
Taurine	1.3 g	0.13 g
Vitamin A	36194.4 IU	3500 IU
Vitamin E	124.1 mg	12 mg
Omega 6	19.6 g	1.9 g
Omega 3	2.4 g	0.23 g
EPA + DHA	1.2 g	0.12 g
Metabolisable energy*		97 kcal / 100 g

\*Calculated according to NRC2006.

### COMPOSITION

Meat and animal derivatives, cereals, oils and fats, derivatives of vegetable origin, minerals, various sugars. Selected source of protein: chicken (38%). Selected source of carbohydrate: rice (4%).

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 350 IU, E1 (Iron): 11 mg, E2 (Iodine): 0.4 mg, E4 (Copper): 3.5 mg, E5 (Manganese): 3.3 mg, E6 (Zinc): 33 mg - Technological additives: Clinoptilolite of sedimentary origin: 0.45 g.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

## SENSITIVITY CONTROL

### CHICKEN WITH RICE

#### WEIGHT MANAGEMENT

#### GASTROINTESTINAL TRACT

#### VITAL SUPPORT

#### HEALTH MANAGEMENT

## CANINE

## SKIN & COAT



Available in 400 g, 1.5 kg, 3.5 kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT CAT	CAT'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	cups	grams	cups	grams	cups
	2	39	3/8	33	3/8	26	2/8
	2.5	46	4/8	38	3/8	30	3/8
	3	52	5/8	43	4/8	35	3/8
	3.5	58	5/8	48	4/8	39	3/8
	4	64	6/8	53	5/8	43	4/8
	4.5	69	6/8	58	5/8	46	4/8
	5	75	7/8	62	5/8	50	4/8
	5.5	80	7/8	67	6/8	53	5/8
	6	85	7/8	71	6/8	57	5/8
	6.5	90	1	75	7/8	60	5/8
	7	95	1	79	7/8	63	6/8
	7.5	100	1+1/8	83	7/8	67	6/8
	8	104	1+1/8	87	1	70	6/8
	8.5	109	1+2/8	91	1	73	6/8
	9	114	1+2/8	95	1	76	7/8
	9.5	118	1+2/8	98	1 1/8	79	7/8
	10	122	1+3/8	102	1 1/8	82	7/8

Feeding instructions: as soon as the first signs of the disease appear, the diet should be prescribed for a 2 month-period to obtain visible results, even though improvement may be noticed as early as 2 weeks after switching diet. For chronic disease, the dermatological diet may be maintained till the cat is 7 years old.

For the mixed feeding table please see the end of the section Dermatology.

## DERMATOSIS & HAIR LOSS



### RECOMMENDED FOR YOUNG\* AND ADULT NEUTERED CATS WITH SENSITIVE SKIN, UP TO 7 YEARS OF AGE

- Feline atopic dermatitis
- Skin and hair support in case of other dermatitis\*\* such as:
  - Flea-bite allergic dermatitis
  - Skin infections or external parasites

\*Post neutering

\*\*In association with medical treatment (such as antiparasitic or antibiotic medications) if needed



#### SKIN BARRIER

Formulated to support the skin's natural protective barrier for optimal health.



#### SKIN & COAT

Contains specific key nutrients, including EPA and DHA, to help support healthy skin and a shiny coat in cats with sensitive skin.



#### LEAN BODY MASS

A high protein diet helps maintain muscle mass.

[S/O®] INDEX

This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	93.8 g	36 g
Fat content	36.5 g	14 g
Carbohydrate	85.9 g	33 g
Crude fibre	9.1 g	3.5 g
Dietary fibre	26.6 g	10.2 g
Calcium	3 g	1.16 g
Phosphorus	2.6 g	1 g
Potassium	2.1 g	0.8 g
Sodium	1.8 g	0.7 g
Chloride	2.4 g	0.93 g
Magnesium	0.2 g	0.07 g
Copper	3.9 mg	1.5 mg
Zinc	44.5 mg	17.1 mg
Taurine	0.6 g	0.23 g
Vitamin A	8333.3 IU	3200 IU
Vitamin E	153.6 mg	59 mg
Omega 6	8.6 g	3.31 g
Omega 3	2.4 g	0.92 g
EPA + DHA	1.1 g	0.43 g
Metabolisable energy*		384 kcal / 100 g

\*Calculated according to NRC2006.

### COMPOSITION

Maize, dehydrated poultry protein, wheat gluten\*\*, rice, maize gluten, hydrolysed animal proteins, animal fats, vegetable fibres, beet pulp, minerals, fish oil, psyllium husks and seeds, fructo-oligo-saccharides, soya oil, borage oil, marigold extract (source of lutein).

\*\*L.I.P.: protein selected for its very high digestibility

### ADDITIVES (KG)\*\*\*

Nutritional additives: Vitamin A: 29500 IU, Vitamin D3: 800 IU, E1 (Iron): 33 mg, E2 (Iodine): 3.4 mg, E4 (Copper): 10 mg, E5 (Manganese): 44 mg, E6 (Zinc): 131 mg, E8 (Selenium): 0.07 mg - Preservatives - Antioxidants.

\*\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

## SKIN & COAT



Available in box of 12 pouches of 85 g pouch

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT CAT	CAT'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	pouch	grams	pouch	grams	pouch
	2	155	2	130	1 + 1/2	105	1
	2.5	180	2	150	2	120	1 + 1/2
	3	205	2 + 1/2	170	2	135	1 + 1/2
	3.5	230	2 + 1/2	190	2	155	2
	4	255	3	210	2 + 1/2	170	2
	4.5	275	3	230	2 + 1/2	185	2
	5	295	3 + 1/2	245	3	200	2 + 1/2
	5.5	315	3 + 1/2	265	3	210	2 + 1/2
	6	335	4	280	3 + 1/2	225	2 + 1/2
	6.5	355	4	300	3 + 1/2	240	3
	7	375	4 + 1/2	315	3 + 1/2	250	3
	7.5	395	4 + 1/2	330	4	265	3
	8	415	5	345	4	275	3
	8.5	430	5	360	4	290	3 + 1/2
	9	450	5 + 1/2	375	4 + 1/2	300	3 + 1/2
	9.5	470	5 + 1/2	390	4 + 1/2	310	3 + 1/2
	10	485	5 + 1/2	405	5	325	4

Feeding instructions: as soon as the first signs of the disease appear, the diet should be prescribed for a 2 month-period to obtain visible results, even though improvement may be noticed as early as 2 weeks after switching diet. For chronic disease, the dermatological diet may be maintained till the cat is 7 years old.

For the mixed feeding table please see the end of the section Dermatology.

## DERMATOSIS & HAIR LOSS



### RECOMMENDED FOR YOUNG\* AND ADULT NEUTERED CATS WITH SENSITIVE SKIN, UP TO 7 YEARS OF AGE

- Feline atopic dermatitis
- Skin and hair support in case of other dermatitis\*\* such as:
  - Flea-bite allergic dermatitis
  - Skin infections or external parasites

\*Post neutering

\*\*In association with medical treatment (such as antiparasitic or antibiotic medications) if needed



#### SKIN BARRIER

Formulated to support the skin's natural protective barrier for optimal health.



#### SKIN & COAT

Contains specific key nutrients, including EPA and DHA, to help support healthy skin and a shiny coat in cats with sensitive skin.



#### ANTIOXYDANT COMPLEX

A synergistic complex of antioxidants to help neutralise free radicals.

[S/O®] INDEX

This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	106.5 g	9.8 g
Fat content	46.7 g	4.3 g
Carbohydrate	50 g	4.6 g
Crude fibre	14.1 g	1.3 g
Dietary fibre	19.6 g	1.8 g
Calcium	2.9 g	0.27 g
Phosphorus	2.3 g	0.21 g
Potassium	1.8 g	0.17 g
Sodium	1.7 g	0.16 g
Chloride	2.5 g	0.23
Magnesium	0.2 g	0.02 g
Copper	4.8 mg	0.44 mg
Zinc	44.6 mg	4.1 mg
Taurine	1.3 g	0.12 g
Vitamin A	59782.6 IU	5500 IU
Vitamin E	119.6 mg	11 mg
Omega 6	15.2 g	1.4 g
Omega 3	2.2 g	0.2 g
EPA + DHA	1.1 g	0.1 g
Metabolisable energy*		92 kcal / 100 g

\*Calculated according to NRC2006.

### COMPOSITION

Meat and animal derivatives, cereals, fish and fish derivatives, oils and fats, derivatives of vegetable origin, vegetable protein extracts, minerals, various sugars.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 310 IU, E1 (Iron): 9 mg, E2 (Iodine): 0.4 mg, E4 (Copper): 3.2 mg, E5 (Manganese): 2.9 mg, E6 (Zinc): 29 mg - Technological additives: Clinoptilolite of sedimentary origin: 0.2 g.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

## SKIN & COAT

WEIGHT MANAGEMENT

GASTROINTESTINAL TRACT

VITAL SUPPORT

HEALTH MANAGEMENT

CANINE

# CATS MIX FEEDING GUIDELINES



24H

MIX: SKIN&COAT DRY + SKIN&COAT WET


CAT'S WEIGHT (kg)	POUCH	WEIGHT CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)
2	1+	19	2/8	12	1/8	5	0
3	1+	33	3/8	24	2/8	13	1/8
4	1+	46	4/8	35	3/8	21	2/8
5	1+	58	5/8	45	4/8	28	2/8
6	1+	70	6/8	55	5/8	35	3/8
7	1+	81	7/8	64	6/8	42	4/8
8	1+	92	1	73	6/8	48	4/8
9	1+	102	1 + 1/8	82	7/8	54	5/8
10	1+	113	1 + 2/8	90	1	60	5/8

# CATS GROWING FEEDING GUIDELINES



24H

SKIN&COAT CAT DRY ONLY  
OR SKIN&COAT CAT WET ONLY  
OR SKIN&COAT CAT MIX DRY + WET

KITTEN'S AGE in months		ADULT WEIGHT					
		3 kg		4 kg		5 kg	
		DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)
6		59	5/8	70	6/8	83	7/8
7		57	5/8	67	6/8	79	7/8
8		54	5/8	64	6/8	75	7/8
9		50	4/8	61	5/8	71	6/8
10		47	4/8	58	5/8	68	6/8
11		45	4/8	55	5/8	65	6/8
12		43	4/8	53	5/8	62	5/8
		POUCH / DAY		POUCH / DAY		POUCH / DAY	
6		3		3 + 1/2		4	
7		2 + 1/2		3		3 + 1/2	
8		2 + 1/2		3		3 + 1/2	
9		2 + 1/2		3		3 + 1/2	
10		2		2 + 1/2		3	
11		2		2 + 1/2		3	
12		2		2 + 1/2		3	
POUCH 		DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)
6	1+	38	3/8	49	4/8	61	5/8
7	1+	35	3/8	46	4/8	58	5/8
8	1+	32	2/8	43	3/8	54	4/8
9	1+	29	2/8	39	3/8	50	4/8
10	1+	26	2/8	36	3/8	46	4/8
11	1+	24	2/8	34	3/8	43	4/8
12	1+	22	2/8	32	1/4	41	3/8







## SATIETY WEIGHT MANAGEMENT



## WEIGHT MANAGEMENT



### RECOMMENDED FOR WEIGHT LOSS AND MAINTENANCE AFTER WEIGHT LOSS IN CASE OF

- Overweight/Obesity
- Stabilized Diabetes Mellitus with overweight/obesity
- Hyperlipidemia with overweight/obesity

### NOT RECOMMENDED IN CASE OF

- Cats with a history of constipation in response to a high fibre diet
- Growth, gestation/lactation



#### EFFECTIVE WEIGHT MANAGEMENT

Provides safe weight loss and helps avoid weight regain. 97% of cats lost weight in 3 months.



#### BEGGING CONTROL

High natural fiber level keeps cats satisfied between meals. Helps control begging in 82% of cats during weight loss.



#### MUSCLE MASS MAINTENANCE

High protein content helps support healthy weight loss while maintaining muscle mass.



**[S/O®]  
INDEX]** This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

Available in 400 g, 1,5 kg, 3,5kg, 6 kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

IDEAL WEIGHT (kg)	STAGE		
	START	4 WEEKS AFTER*	MAINTENANCE AFTER WEIGHT LOSS
	grams	grams	grams
2	29	23	30
2.5	33	27	35
3	38	30	40
3.5	42	34	45
4	47	37	49
4.5	51	40	54
5	55	43	58
5.5	59	46	62
6	62	49	66
6.5	66	52	70
7	70	55	73
7.5	73	58	77
8	76	61	81

\* If the cat did not lose weight

Individual target weight is determined and then feeding amount established. The recommended daily intake is based on the pet's ideal weight. The starting ration may need to be adjusted during follow-up visits, based on each individual's response and the rate of weight loss. An objective of losing 1-3% bodyweight per week is generally achievable over the first 3 months, and of 0.5-2% afterwards.

**How is the energy allocation for maintenance after weight loss calculated?** Energy requirements to maintain stable body weight after weight loss are lower than those of cats that have never been obese. Based on current scientific insights [Serisier & al. 2011], our maintenance after weight loss rations are calculated by increasing the weight loss ration by +33%.

**NOTE: maintenance rations may need to be adjusted based on individual evolution.** Serisier S. & al. Weight loss and post weight loss maintenance energy requirement of obese colony cats. Proc. 21th ECVIM Congress. Sevilla 2011; 209-210.

KEY VALUES	Per 1000 kcal	Per 100g as fed
Protein	111.84 g	34.0 g
Fat content	29.61 g	9.0 g
Digestible Carbohydrates (Starch)	61.18 g	18.6 g
NFE	93.09 g	28.3 g
Crude fibre	48.03 g	14.6 g
Dietary fibre	79.93 g	24.3 g
Omega 6	6.09 g	1.85 g
Omega 3	1.55 g	0.47 g
EPA + DHA	0.66 g	0.20 g
Calcium	4.44 g	1.35 g
Phosphorus	3.72 g	1.13 g
Sodium	1.81 g	0.55 g
Metabolisable energy*		304 kcal/100 g

\*Calculated according to NRC2006TDF.

### COMPOSITION

Dehydrated poultry protein, vegetable fibres, tapioca, wheat flour, hydrolysed animal proteins, wheat gluten\*, maize gluten, animal fats, chicory pulp, minerals, psyllium husks and seeds, fish oil, marigold extract (source of lutein), glucosamine, hydrolysed cartilage (source of chondroitin).

\*L.I.P. (Low Indigestible Protein): protein selected for its very high digestibility.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin A: 25000 IU, Vitamin D3: 1000 IU, Iron [3b103]: 33 mg, Iodine [3b201, 3b202]: 3.3 mg, Copper [3b405, 3b406]: 10 mg, Manganese [3b502, 3b504]: 43 mg, Zinc [3b603, 3b605, 3b606]: 133 mg, Selenium [3b801, 3b811, 3b812]: 0.05 mg - Preservatives - Antioxidants.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

## SATIETY WEIGHT MANAGEMENT



## WEIGHT MANAGEMENT



### RECOMMENDED FOR WEIGHT LOSS AND MAINTENANCE AFTER WEIGHT LOSS IN CASE OF

- Overweight/Obesity
- Stabilized Diabetes Mellitus with overweight/obesity
- Hyperlipidemia with overweight/obesity

### NOT RECOMMENDED IN CASE OF

- Growth, gestation/lactation



#### EFFECTIVE WEIGHT MANAGEMENT

Helps provide safe weight loss and avoid weight regain.



#### BEGGING CONTROL

Nutritional concept that helps keeping cats satisfied between meals



#### MUSCLE MASS MAINTENANCE

High protein content helps maintain muscle mass during weight loss



**[S/O®]  
INDEX]** This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

Available in box of 12 pouches of 85 g

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

IDEAL WEIGHT (kg)	STAGE					
	START		4 WEEKS AFTER*		MAINTENANCE AFTER WEIGHT LOSS	
	grams	pouch	grams	pouch	grams	pouch
2	130	1 + 1/2	105	1	140	1 + 1/2
2.5	155	2	120	1 + 1/2	165	2
3	175	2	140	1 + 1/2	185	2
3.5	195	2 + 1/2	155	2	205	2 + 1/2
4	215	2 + 1/2	170	2	225	2 + 1/2
4.5	235	3	185	2	245	3
5	250	3	200	2 + 1/2	265	3
5.5	270	3	215	2 + 1/2	285	3 + 1/2
6	285	3 + 1/2	225	2 + 1/2	305	3 + 1/2
6.5	305	3 + 1/2	240	3	320	4
7	320	4	255	3	340	4
7.5	335	4	265	3	355	4
8	350	4	280	3 + 1/2	370	4 + 1/2

\* If the cat did not lose weight

Individual target weight is determined and then feeding amount established. The recommended daily intake is based on the pet's ideal weight. The starting ration may need to be adjusted during follow-up visits, based on each individual's response and the rate of weight loss. An objective of losing 1-3% bodyweight per week is generally achievable over the first 3 months, and of 0.5-2% afterwards.

**How is the energy allocation for maintenance after weight loss calculated?** Energy requirements to maintain stable body weight after weight loss are lower than those of cats that have never been obese.

Based on current scientific insights [Serisier & al. 2011], our maintenance after weight loss rations are calculated by increasing the weight loss ration by +33%.

**NOTE: maintenance rations may need to be adjusted based on individual evolution.** Serisier S. & al. Weight loss and post weight loss maintenance energy requirement of obese colony cats. Proc. 21th ECVIM Congress. Sevilla 2011; 209-210.

KEY VALUES	Per 1000 kcal	Per 100g as fed
Protein	113.64 g	7.5 g
Fat content	33.33 g	2.2 g
Digestible Carbohydrates (Starch)	33.33 g	2.2 g
NFE	50.00 g	3.3 g
Crude fibre	19.70 g	1.3 g
Dietary fibre	22.73 g	1.5 g
Omega 6	7.58 g	0.50 g
Omega 3	0.76 g	0.05 g
EPA + DHA	0.06 g	0.00 g
Calcium	4.09 g	0.27 g
Phosphorus	3.33 g	0.22 g
Sodium	2.42 g	0.16 g
Metabolisable energy*		66 kcal/100 g

\*Calculated according to NRC2006TDF.

### COMPOSITION

Meat and animal derivatives, cereals, derivatives of vegetable origin, eggs and egg derivatives, minerals, various sugars, molluscs and crustaceans.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 235 IU, Iron [3b103]: 7 mg, Iodine [3b202]: 0.23 mg, Copper [3b405, 3b406]: 1.9 mg, Manganese [3b502, 3b503, 3b504]: 2 mg, Zinc [3b603, 3b605, 3b606]: 20 mg.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

SATIETY WEIGHT MANAGEMENT



SATIETY WEIGHT MANAGEMENT



GASTROINTESTINAL TRACT

VITAL SUPPORT

HEALTH MANAGEMENT

CANINE

# DIABETIC



Available in 400 g, 1.5 kg, 3.5 kg bags

## RECOMMENDED DAILY INTAKE & FEEDING DURATION

CAT'S WEIGHT (kg)	BODY CONDITION					
	THIN		NORMAL		OVERWEIGHT	
	grams	cups	grams	cups	grams	cups
2	40	4/8	33	3/8	26	2/8
2.5	46	4/8	39	3/8	31	3/8
3	53	5/8	44	4/8	35	3/8
3.5	59	5/8	49	4/8	39	3/8
4	65	6/8	54	5/8	43	4/8
4.5	70	6/8	59	5/8	47	4/8
5	76	7/8	63	6/8	51	4/8
5.5	81	7/8	68	6/8	54	5/8
6	86	1	72	6/8	58	5/8
6.5	91	1	76	7/8	61	5/8
7	96	1	80	7/8	64	6/8
7.5	101	1 + 1/8	84	7/8	67	6/8
8	106	1 + 1/8	88	1	71	6/8
8.5	111	1 + 2/8	92	1	74	6/8
9	115	1 + 2/8	96	1	77	7/8
9.5	120	1 + 3/8	100	1 + 1/8	80	7/8
10	124	1 + 3/8	104	1 + 1/8	83	7/8

DIABETIC is recommended as first line nutritional management for all newly diagnosed cases of Diabetes Mellitus until control of clinical signs and hyperglycemia have been achieved.

NOTE: the use of this food may affect the cat's sensitivity to insulin. It is highly recommended that blood glucose and insulin response are monitored when starting the diet so that insulin dosage can be adapted. Once glycemic control is reached, nutritional management should be guided by assessment of the Body Condition Score.

BCS 4 or less: continue DIABETIC  
BCS 7-8-9: transition to SATIETY WEIGHT MANAGEMENT

It is highly recommended that blood glucose and insulin response are monitored when starting or changing the diet so that insulin dosage can be adapted.

# DIABETES MELLITUS




## RECOMMENDED FOR CASES OF

- Diabetes Mellitus and a Body Condition Score (BCS) ≤ 6/9
- Diabetes Mellitus and a BCS ≥ 7/9, until glycemic control has been achieved

## NOT RECOMMENDED IN CASE OF

- Growth, gestation/lactation



### GLUCOMODULATION

Specific formula to help in the management of post-prandial blood glucose in diabetic cats.



### HIGH PROTEIN

High protein content. Maintenance of muscle mass is essential in diabetic cats.



### LOW STARCH

Formula that contains a reduced level of starch.

[S/O®] This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

## KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	119.42 g	46.0 g
Fat content	31.15 g	12.0 g
Digestible Carbohydrates (Starch)	48.81 g	18.8 g
NFE	68.28 g	26.3 g
Crude fibre	9.87 g	3.8 g
Dietary fibre	29.34 g	11.3 g
Omega 6	7.14 g	2.75 g
Omega 3	1.87 g	0.72 g
EPA + DHA	0.78 g	0.30 g
Calcium	2.39 g	0.92 g
Phosphorus	2.36 g	0.91 g
Sodium	1.04 g	0.40 g
Arginine	7.09 g	2.73 g
Metabolisable energy*		385 kcal/100 g

\*Calculated according to NRC2006CF.

## COMPOSITION

Dehydrated poultry protein, barley, wheat gluten\*, maize gluten, soya protein isolate\*, tapioca, animal fats, hydrolysed animal proteins, vegetable fibres, chicory pulp, minerals, fish oil, psyllium husks and seeds, fructo-oligo-saccharides, soya oil, marigold extract (source of lutein). Sources of carbohydrates: barley, tapioca.

\*L.I.P. (Low Indigestible Protein): protein selected because of its very high assimilation.

## ADDITIVES (KG)\*\*

Nutritional additives: Vitamin A: 25000 IU, Vitamin D3: 1000 IU, Iron (3b103): 33 mg, Iodine (3b201, 3b202): 3.3 mg, Copper (3b405, 3b406): 10 mg, Manganese (3b502, 3b504): 43 mg, Zinc (3b603, 3b605, 3b606): 130 mg, Selenium (3b801, 3b811, 3b812): 0.05 mg - Preservatives - Antioxidants.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



# DIABETIC



# DIABETIC



Available in box of 12 pouches of 85 g

## RECOMMENDED DAILY INTAKE & FEEDING DURATION

CAT'S WEIGHT (kg)	BODY CONDITION					
	THIN		NORMAL		OVERWEIGHT	
	grams	pouch	grams	pouch	grams	pouch
2	195	2 + 1/2	165	2	130	1 + 1/2
2.5	230	2 + 1/2	190	2	155	2
3	260	3	220	2 + 1/2	175	2
3.5	290	3 + 1/2	245	3	195	2 + 1/2
4	320	4	270	3	215	2 + 1/2
4.5	350	4	290	3 + 1/2	235	3
5	375	4 + 1/2	315	3 + 1/2	250	3
5.5	405	5	335	4	270	3
6	430	5	355	4	285	3 + 1/2
6.5	455	5 + 1/2	380	4 + 1/2	300	3 + 1/2
7	480	5 + 1/2	400	4 + 1/2	320	4
7.5	500	6	420	5	335	4
8	525	6	440	5	350	4
8.5	550	6 + 1/2	460	5 + 1/2	365	4 + 1/2
9	570	6 + 1/2	475	5 + 1/2	380	4 + 1/2
9.5	595	7	495	6	395	4 + 1/2
10	615	7	515	6	410	5

It is recommended to feed DIABETIC for the cat's lifetime.

NOTE: The use of this food may affect the cat's sensitivity to insulin. It is highly recommended that blood glucose and insulin response are monitored when starting the diet so that insulin dosage can be adapted.

# DIABETES MELLITUS




## RECOMMENDED FOR CASES OF

- Diabetes Mellitus and a Body Condition Score (BCS) ≤ 6/9
- Diabetes Mellitus and a BCS ≥ 7/9, until glycemic control has been achieved

## NOT RECOMMENDED IN CASE OF

- Growth, gestation/lactation



### GLUCOMODULATION

Specific formula to help in the management of post-prandial blood glucose in diabetic cats.



### HIGH PROTEIN

High protein content. Maintenance of muscle mass is essential in diabetic cats.



### LOW STARCH

Formula that contains a reduced level of starch.

[S/O®] This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

## KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	114.69 g	8.9 g
Fat content	41.24 g	3.2 g
Digestible Carbohydrates (Starch)	15.46 g	1.2 g
NFE	33.51 g	2.6 g
Crude fibre	14.18 g	1.1 g
Dietary fibre	19.33 g	1.5 g
Omega 6	7.73 g	0.60 g
Omega 3	2.19 g	0.17 g
EPA + DHA	1.38 g	0.11 g
Calcium	3.74 g	0.29 g
Phosphorus	2.71 g	0.21 g
Sodium	2.06 g	0.16 g
Arginine	7.09 g	0.55 g
Metabolisable energy*		78 kcal/100 g

\*Calculated according to NRC2006TDF.

## COMPOSITION

Meat and animal derivatives, derivatives of vegetable origin, cereals, minerals, oils and fats.

## ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 270 IU, Iron (3b103): 3 mg, Iodine (3b201, 3b202): 0.4 mg, Copper (3b405, 3b406): 3 mg, Manganese (3b502, 3b504): 0.8 mg, Zinc (3b603, 3b605, 3b606): 8 mg - Technological additives: Clinoptilolite of sedimentary origin: 1 g.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



# DIABETIC



# CATS MIX FEEDING GUIDELINES



24H

MIX: SATIETY CAT DRY + SATIETY CAT WET POUCH

IDEAL WEIGHT (kg)	POUCH	STAGE		
		START	4 WEEKS AFTER*	MAINTENANCE AFTER WEIGHT LOSS
		DRY (g)	DRY (g)	DRY (g)
2	1+	10	5	13
3	1+	20	13	24
4	1+	28	21	34
5	1+	36	28	43
6	1+	44	35	52
7	1+	51	41	61
8	1+	58	47	69

\* If the cat did not lose weight



24H

MIX: DIABETIC CAT DRY + DIABETIC CAT WET POUCH

CAT'S WEIGHT (kg)	POUCH	WEIGHT CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)
2	1+	23	2/8	17	1/8	9	1/8
3	1+	38	3/8	29	3/8	18	2/8
4	1+	51	4/8	40	3/8	26	2/8
5	1+	64	6/8	50	4/8	33	3/8
6	1+	75	7/8	60	5/8	40	4/8
7	1+	87	1	70	6/8	47	4/8
8	1+	98	1 + 1/8	79	7/8	53	5/8
9	1+	108	1 + 2/8	88	1	60	5/8
10	1+	119	1 + 2/8	96	1	66	6/8



# CANINE

VETERINARY HEALTH NUTRITION

CANINE







# INDEX OF DOG HEALTH ISSUES

PLEASE NOTE: All dietary recommendations throughout this book are for the nutritional management of cats exhibiting listed conditions and do not preclude appropriate medical management. All clinical diets should be used under veterinary supervision.

A	Adverse Food Reactions (AFR: Food allergy/Food intolerance)			-HYPOALLERGENIC canine
				-SENSITIVITY CONTROL canine
				-ANALLERGENIC canine
				-MULTIFUNCTION RENAL-HYPOALLERGENIC concomitant with Chronic Kidney Disease (CKD)
	Anorexia	Poor appetite		-GASTRO-INTESTINAL canine
		Tube feeding		-CONVALESCENCE SUPPORT INSTANT
				-RECOVERY
	Ascites	With heart failure		-CARDIAC canine
		With hypoproteinemia	Lymphangiectasia (protein losing enteropathy)	-GASTRO-INTESTINAL canine Low FAT
			Liver disease	-HEPATIC canine
Protein losing nephropathy			-RENAL canine	
Atopy (CAD: Canine atopic dermatitis)				see dermatitis
C	Cachexia			-CONVALESCENCE SUPPORT INSTANT
				-RECOVERY
				-GASTRO-INTESTINAL canine (EXCEPT MODERATE CALORIE OR LOW FAT)
	Cancer			see neoplasia
	Chronic Kidney Disease (CKD)			-RENAL canine
	Concomitant with Adverse Food reactions			-MULTIFUNCTION canine GFR-AFR
	Early CKD (IRIS stage 1)			-VCN SENIOR CONSULT MATURE canine
	Azotemic CKD (IRIS stages 2 to 4)			-RENAL canine
	Colitis			see diarrhea
	Constipation			-GASTRO-INTESTINAL canine
	Convalescence			-RECOVERY
				-CONVALESCENCE SUPPORT INSTANT
				-GASTRO-INTESTINAL canine (except moderate calorie or low fat)
	Cushing's syndrome (hyperadrenocorticism)			No specific therapeutic diet, it depends on patient and associated conditions
	With Diabetes Mellitus			-DIABETIC canine
	With chronic kidney disease			-RENAL canine
Cystitis				see Urinary Tract Infection
D	Dermatitis	Atopic dermatitis		-SKIN SUPPORT canine
				-SKIN CARE canine
		Associated with Adverse Food Reaction	-HYPOALLERGENIC canine	
			-SENSITIVITY CONTROL canine	
			-ANALLERGENIC canine	
		Cutaneous Adverse Food Reactions (CAFR)	-HYPOALLERGENIC canine	
			-SENSITIVITY CONTROL canine	
			-ANALLERGENIC canine	
		Concomitant with Chronic Kidney Disease	-MULTIFUNCTION RENAL + HYPOALLERGENIC canine	
		Concomitant with Struvite or Calcium Oxalate urolithiasis	-MULTIFUNCTION URINARY + HYPOALLERGENIC canine	
		Flea Bite Allergic Dermatitis (in adjunction to flea treatment)	-SKIN SUPPORT canine	
			-SKIN CARE canine	
		Pyodermatitis (in adjunction to adequate antibiotic therapy)	-SKIN SUPPORT canine	
			-SKIN CARE canine	



PLEASE NOTE: The following are recommendations only. The veterinary diets should be used under veterinary supervision.

D	Diabetes Mellitus (in adjunction to insulin therapy)	During stabilization phase (until control of clinical signs and hyperglycemia) and for patients with BCS ≤6/9	-DIABETIC canine
		For stabilized patients with BCS ≥7/9	-SATIETY canine -SATIETY SMALL DOG canine
		For patients with BCS ≥7/9 and concomitant struvite or calcium oxalate urolithiasis	-URINARY S/O SATIETY canine
	Diarrhea	Acute diarrhea	-GASTRO-INTESTINAL canine
		Non-specific or unknown aetiology:	-GASTRO-INTESTINAL canine
		Group of chronic enteropathies/ Inflammatory Bowel Disease	-HYPOALLERGENIC canine -SENSITIVITY CONTROL canine -ANALLERGENIC canine
		Food responsive diarrhea	-GASTRO-INTESTINAL canine
		Antibiotic responsive diarrhea (in adjunction to adequate antibiotherapy)	-GASTRO-INTESTINAL canine
		Immunosuppressive drug responsive diarrhea (in adjunction to anti-inflammatories/ immunosuppressors)	-GASTRO-INTESTINAL canine -HYPOALLERGENIC canine -SENSITIVITY CONTROL canine -ANALLERGENIC canine
		Lymphangiectasia (protein-losing enteropathy)	-GASTRO-INTESTINAL canine LOW FAT
		Small intestinal bacterial overgrowth (SIBO)	-GASTRO-INTESTINAL canine
		Fibre responsive colitis (including stress diarrhea)	-FIBRE RESPONSE canine
		Chronic diarrhea	
E	Enteropathies (chronic enteropathies)		see diarrhoea (chronic diarrhoea)
	Exocrine pancreatic insufficiency (EPI) (in adjunction to digestive enzyme treatment)		-GASTRO-INTESTINAL canine -HYPOALLERGENIC canine
F	Food allergies/ Food intolerances		see Adverse Food Reactions
	Food elimination trial		-ANALLERGENIC canine -HYPOALLERGENIC canine -SENSITIVITY CONTROL canine
G	Gastric disorders	Delayed gastric emptying	-GASTRO-INTESTINAL canine LOW FAT
		Gastritis, gastric and/or duodenal ulcer	-HYPOALLERGENIC canine -SENSITIVITY CONTROL canine -GASTRO-INTESTINAL canine
	Gastro-enteritis		see diarrhea
H	Heart disease/failure		-CARDIAC canine
	Hepatic disease	Hepatitis	-GASTRO-INTESTINAL canine MODERATE CALORIE
		Acute hepatitis	-RECOVERY
		Chronic hepatitis	-HEPATIC canine
		Copper storage disease	-HEPATIC canine
		Portosystemic shunt	-HEPATIC canine
		Hepatic encephalopathy	-HEPATIC canine
	Hyperadrenocorticism		see Cushing's syndrome
	Hyperlipidemia (fasting hyperlipidemia)	Severe	-GASTRO-INTESTINAL canine LOW FAT
		Moderate	-GASTRO-INTESTINAL canine -MODERATE CALORIE canine
	Hypertension		-CARDIAC canine
	Hypothyroidism		No specific therapeutic diet, focus on associated signs
		If hyperlipidemia	-GASTRO-INTESTINAL canine LOW FAT
		If overweight	-SATIETY canine -SATIETY SMALL DOG canine
I	Inflammatory bowel disease		see Diarrhea (chronic diarrhoea)
K	Kidney disease		see CKD (Chronic Kidney Disease)
L	Leishmaniasis	Prevention of urate urolithiasis secondary to allopurinol treatment	-URINARY U/C LOW PURINE canine
		If secondary renal disease (glomerulonephritis)	-RENAL canine
	Liver disease		see Hepatic disease
	Lymphangiectasia		see Diarrhea (chronic diarrhoea)


PLEASE NOTE: The following are recommendations only. The veterinary diets should be used under veterinary supervision.

M	Maldigestion / Malabsorption		-GASTRO-INTESTINAL canine -HYPOALLERGENIC canine -ANALLERGENIC canine
	Malnutrition		see Cachexia
	Megacolon		-GASTRO-INTESTINAL canine
	Megaesophagus	Gastrostomy feeding tube	-CONVALESCENCE SUPPORT INSTANT -RECOVERY -GASTRO-INTESTINAL HIGH ENERGY canine LIQUID
		Oral feeding (depending on individual response to liquid, gruel or balls form)	-CONVALESCENCE SUPPORT INSTANT -RECOVERY -GASTRO-INTESTINAL canine -GASTRO-INTESTINAL HIGH ENERGY canine LIQUID
N	Neoplasia (optimal nutrition during cancer treatment)		-RECOVERY -GASTRO-INTESTINAL canine (except Moderate calorie or Low fat)
O	Obesity/ Overweight (weight loss and weight maintenance after weight loss)		-SATIETY canine
		Associated with urolithiasis	-MULTIFUNCTION URINARY + SATIETY canine
		Associated with osteo-articular disorders	-MULTIFUNCTION MOBILITY C2P+ + SATIETY canine
	Oesophagitis	Associated with delayed gastric emptying / reflux	-GASTRO-INTESTINAL canine LOW FAT WET (OR KIBBLES SOAKED IN WATER*)
		Other causes	-RECOVERY
	Oral hygiene		-DENTAL canine
	Osteo-articular disorders		-MOBILITY C2P+ canine
P	Otitis (Atopic Dermatitis and/or Adverse Food Reaction with otitis externa)		-SKIN SUPPORT canine -SKIN CARE canine -HYPOALLERGENIC canine
	Pancreatitis	Acute pancreatitis	-GASTRO-INTESTINAL canine LOW FAT
		Chronic pancreatitis	-GASTRO-INTESTINAL canine LOW FAT
		Severe	-GASTRO-INTESTINAL canine MODERATE CALORIE
		Moderate	-HYPOALLERGENIC canine MODERATE CALORIE -SENSITIVITY CONTROL canine DRY
		If associated with obesity/overweight	-SATIETY canine
R	Perianal fistula	If complicated with diabetes	-DIABETIC canine
			-HYPOALLERGENIC canine
			-ANALLERGENIC canine
	Poor appetite		-RECOVERY -GASTRO-INTESTINAL canine (except Moderate calorie or Low fat)
	Pre/post-surgery: (including digestive surgery)		-RECOVERY -CONVALESCENCE SUPPORT INSTANT -GASTRO-INTESTINAL canine
	Protein-losing enteropathy		see Diarrhea (chronic diarrhoea)
	Renal disease/ failure		see CKD (Chronic Kidney Disease)
		Acute Kidney Injury	-RENAL LIQUID canine
S	Small intestinal bacterial overgrowth (SIBO)		see Diarrhea (chronic diarrhoea)
	Stress-and anxiety-related signs		-CALM canine



PLEASE NOTE: The following are recommendations only. The veterinary diets should be used under veterinary supervision.

T	Tartar and dental plaque			-DENTAL canine
	Tube feeding	Tube with diameter <10Fr	Need of protein and/or phosphorus restriction (kidney disease, hepatic encephalopathy...)	-RENAL LIQUID canine
			Need of fat restriction (pancreatitis, lymphangiectasia, hyperlipidemia...)	- GASTRO- INTESTINAL LOW FAT LIQUID canine
			Need of higher energy/digestive support	- GASTRO- INTESTINAL HIGH ENERGY LIQUID canine
			All other conditions	-RECOVERY LIQUID / CONVALESCENCE SUPPORT INSTANT canine
		Tube with diameter >10Fr	Non-specific conditions	-RECOVERY LIQUID OR WET / CONVALESCENCE SUPPORT INSTANT canine
			Need of protein and/or phosphorus restriction (kidney disease, hepatic encephalopathy...)	-RENAL LIQUID canine
			Need of fat restriction (pancreatitis, lymphangiectasia, hyperlipidemia...)	- GASTRO- INTESTINAL LOW FAT LIQUID canine
			Need of higher energy/digestive support	- GASTRO- INTESTINAL HIGH ENERGY LIQUID canine
		Other specific conditions: corresponding specific diet mixed with water and blended (wet diet with same volume of water,or kibbles soaked with water*)		
	U	Urolithiasis	Struvite urolithiasis (prevention & dissolution)	-URINARY S/O canine
		Associated with Adverse Food Reactions	-MULTIFUNCTION URINARY HYPOALLERGENIC canine	
		Associated with overweight/obesity	-MULTIFUNCTION URINARY + SATIETY canine	
		Calcium oxalate or calcium phosphate urolithiasis (prevention)	-URINARY S/O canine	
		Associated with Adverse Food Reactions	-MULTIFUNCTION URINARY HYPOALLERGENIC canine	
		Associated with overweight/obesity	-MULTIFUNCTION URINARY + SATIETY canine	
		Purine urolithiasis (urate, xanthine) (prevention)	-URINARY U/C LOW PURINE canine DRY, KIBBLES SOAKED WITH WATER*	
		Urate stones associated with porto-systemic shunt	- HEPATIC canine DRY, KIBBLES SOAKED WITH WATER*	
		Purine urolithiasis associated with Adverse Food Reactions	-ANALLERGENIC canine DRY, KIBBLES SOAKED WITH WATER*	
		Cystine urolithiasis (prevention)	-URINARY U/C LOW PURINE canine DRY, KIBBLES SOAKED WITH WATER*	
			-RENAL canine WET CHUNKS IN GRAVY	
	Urinary Tract Infection (UTI) (dissolution of struvite concomitant with UTI) (in adjunction to adequate antibiotherapy)			-URINARY S/O canine



\* Kibbles rehydration for blending  
- mix one volume of kibbles with 3 volumes of hot tap water (around 60°)  
- let the kibbles absorb water for about 15 min before serving to the dog  
(you can blend it for smoother texture)

\*\* Increasing water consumption in cats (cats being less prone than dogs to consume rehydrated kibbles) Water-related factors to consider include freshness, taste, movement (water fountains, dripping faucets, bowl with pump-bubbled air...), and shape of the container.





## URINARY S/O



Available in 2 kg, 7.5 kg, 13 kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

DOG'S WEIGHT (kg)	BODY CONDITION					
	THIN		NORMAL		OVERWEIGHT	
	grams	cups	grams	cups	grams	cups
2	54	5/8	48	4/8	41	4/8
4	91	1	80	7/8	69	6/8
6	123	1 + 3/8	109	1 + 2/8	94	1 + 1/8
8	153	1 + 6/8	135	1 + 5/8	116	1 + 3/8
10	181	2 + 1/8	159	1 + 7/8	138	1 + 5/8
15	245	2 + 7/8	216	2 + 4/8	187	2 + 1/8
20	305	3 + 4/8	268	3 + 1/8	232	2 + 5/8
25	360	4 + 1/8	317	3 + 5/8	274	3 + 1/8
30	413	4 + 6/8	363	4 + 2/8	314	3 + 5/8
35	463	5 + 3/8	408	4 + 6/8	352	4 + 1/8
40	512	5 + 7/8	451	5 + 2/8	389	4 + 4/8
45	560	6 + 4/8	492	5 + 6/8	425	4 + 7/8
50	606	7	533	6 + 1/8	460	5 + 3/8
55	650	7 + 4/8	572	6 + 5/8	494	5 + 6/8
60	694	8	611	7 + 1/8	528	6 + 1/8
70	779	9	686	8	592	6 + 7/8
80	862	10	758	8 + 6/8	655	7 + 5/8

Urinary S/O is a complete dietetic feed for dogs, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties, its low level of magnesium and its restricted level of protein but of high quality.

RECOMMENDATIONS: It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. Then the dog should be re-evaluated regularly by the veterinarian to assess the relevance of a life long recommendation. Water should be available at all times.

NOTE: - renal function should be checked before the advice of a new diet.  
- in older dogs, the use of Urinary S/O Ageing\* is recommended.

## LOWER URINARY TRACT DISEASE



### RECOMMENDED FOR CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotherapy)

### NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying drugs
- Growth, gestation/lactation



#### STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



#### LOW RSS

Helps lower the concentration of ions contributing to crystal formation.



#### URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



#### LOW MAGNESIUM

Reduced level of magnesium, a natural component of struvite crystals.



This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	46.38 g	18 g
Fat content	43.8 g	17 g
Carbohydrate	120.85 g	46.9 g
Crude fibre	5.67 g	2.2 g
Dietary fibre	14.69 g	5.7 g
Calcium	1.29 g	0.5 g
Phosphorus	1.29 g	0.5 g
Potassium	2.06 g	0.8 g
Sodium	2.83 g	1.1 g
Chloride	5.51 g	2.14 g
Magnesium	0.1 g	0.04 g
Copper	3.86 mg	1.5 mg
Zinc	43.8 mg	17 mg
Taurine	0.54 g	0.21 g
Vitamin A	4380.31 IU	1700 IU
Vitamin E	157.18 mg	61 mg
Omega 6	9.25 g	3.59 g
Omega 3	1.67 g	0.63 g
EPA + DHA	0.54 g	0.21 g
Metabolisable energy*		388 kcal / 100 g

\*Calculated according to NRC2006

### COMPOSITION

Maize flour, rice, animal fats, dehydrated poultry protein, maize gluten, minerals, hydrolysed animal proteins, vegetable fibres, soya oil, fish oil, fructo-oligo-saccharides, mono - and diglycerides of palmitic and stearic acids esterified with citric acid, marigold extract (source of lutein). Protein sources: dehydrated poultry protein, hydrolysed animal proteins.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin A: 15500 IU, Vitamin D3: 1000 IU, E1 (Iron): 44 mg, E2 (Iodine): 4.4 mg, E4 (Copper): 13 mg, E5 (Manganese): 57 mg, E6 (Zinc): 147 mg, E8 (Selenium): 0.06 mg. Urine acidifying substances: calcium sulfate (0.88%). Preservatives - Antioxidants.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



## URINARY S/O



## URINARY S/O



LOAF



Available in 200 g can

### RECOMMENDED DAILY INTAKE & FEEDING DURATION BASED ON 200 G

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	can	grams	can	grams	can
	2	208	1	183	1	158	3/4
	4	350	1 + 3/4	308	1 + 2/4	266	1 + 1/4
	6	475	2 + 2/4	418	2	361	1 + 3/4
	8	589	3	519	2 + 2/4	448	2 + 1/4
	10	697	3 + 2/4	613	3	529	2 + 3/4
	15	944	4 + 3/4	831	4 + 1/4	718	3 + 2/4
	20	1 172	5 + 3/4	1 031	5 + 1/4	890	4 + 2/4
	25	1 385	7	1 219	6	1 053	5 + 1/4
	30	1 588	8	1 397	7	1 207	6
	35	1 783	9	1 569	7 + 3/4	1 355	6 + 3/4
	40	1 970	9 + 3/4	1 734	8 + 3/4	1 498	7 + 2/4
	45	2 152	10 + 3/4	1 894	9 + 2/4	1 636	8 + 1/4
	50	2 329	11 + 3/4	2 050	10 + 1/4	1 770	8 + 3/4
	55	2 502	12 + 2/4	2 202	11	1 902	9 + 2/4
	60	2 671	13 + 1/4	2 350	11 + 3/4	2 030	10 + 1/4
	70	2 998	15	2 638	13 + 1/4	2 279	11 + 2/4
	80	3 314	16 + 2/4	2 916	14 + 2/4	2 519	12 + 2/4

Urinary S/O is a complete dietetic feed for dogs, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties, its low level of magnesium and its restricted level of protein but of high quality.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. Then the dog should be re-evaluated regularly by the veterinarian to assess the relevance of a life long recommendation. Water should be available at all times.

**NOTE:** -renal function should be checked before the advice of a new diet.  
-in older dogs, the use of Urinary S/O Ageing™ is recommended.

## LOWER URINARY TRACT DISEASE



### RECOMMENDED FOR CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotherapy)

### NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying drugs
- Growth, gestation/lactation



#### STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



#### URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



#### LOW RSS

Helps lower the concentration of ions contributing to crystal formation.

**[S/O®] INDEX** This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	83.68 g	8 g
Fat content	62.76 g	6 g
Carbohydrate	48.12 g	4.6 g
Crude fibre	14.64 g	1.4 g
Dietary fibre	25.1 g	2.4 g
Calcium	2.2 g	0.21 g
Phosphorus	2.09 g	0.2 g
Potassium	2.09 g	0.2 g
Sodium	3.03 g	0.29 g
Chloride	4.18 g	0.4 g
Magnesium	0.17 g	0.02 g
Copper	3.14 mg	0.3 mg
Zinc	23.01 mg	2.2 mg
Taurine	1.26 g	0.12 g
Vitamin A	47071.13 IU	4500 IU
Vitamin E	198.74 mg	19 mg
Omega 6	25.1 g	2.4 g
Omega 3	1.36 g	0.13 g
EPA + DHA	0.63 g	0.06 g
Metabolisable energy*		96 kcal / 100 g

\*Calculated according to NRC2006.

### COMPOSITION

Meat and animal derivatives, oils and fats, cereals, derivatives of vegetable origin, minerals, various sugars.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 190 IU, E1 (Iron): 3.4 mg, E2 (Iodine): 0.24 mg, E4 (Copper): 2 mg, E5 (Manganese): 1 mg, E6 (Zinc): 10 mg, Urine acidifying substances: calcium sulfate: 0.25%.

Technological additives: Sodium bisulfate: 0.56 g.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



## URINARY S/O



## URINARY S/O



LOAF



Available in 410 g can

### RECOMMENDED DAILY INTAKE & FEEDING DURATION BASED ON 410 G

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	can	grams	can	grams	can
	2	208	1/2	183	1/2	158	1/2
	4	350	3/4	308	3/4	266	3/4
	6	475	1 + 1/4	418	1	361	1
	8	589	1 + 1/2	519	1 + 1/4	448	1
	10	697	1 + 3/4	613	1 + 1/2	529	1 + 1/4
	15	944	2 + 1/4	831	2	718	1 + 3/4
	20	1 172	1 + 3/4	1 031	2 + 1/2	890	2 + 1/4
	25	1 385	3 + 1/2	1 219	3	1 053	2 + 1/2
	30	1 588	3 + 3/4	1 397	3 + 1/2	1 207	3
	35	1 783	4 + 1/4	1 569	3 + 3/4	1 355	3 + 1/4
	40	1 970	4 + 3/4	1 734	4 + 1/4	1 498	3 + 3/4
	45	2 152	5 + 1/4	1 894	4 + 1/2	1 636	4
	50	2 329	5 + 3/4	2 050	5	1 770	4 + 1/4
	55	2 502	6	2 202	5 + 1/4	1 902	4 + 3/4
	60	2 671	6 + 1/2	2 350	5 + 3/4	2 030	5
	70	2 998	7 + 1/4	2 638	6 + 1/2	2 279	5 + 1/2
	80	3 314	8	2 916	7	2 519	6 + 1/4

Urinary S/O is a complete dietetic feed for dogs, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties, its low level of magnesium and its restricted level of protein but of high quality.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. Then the dog should be re-evaluated regularly by the veterinarian to assess the relevance of a life long recommendation. Water should be available at all times.

**NOTE:** -renal function should be checked before the advice of a new diet.  
-in older dogs, the use of Urinary S/O Ageing™ is recommended.

## LOWER URINARY TRACT DISEASE



### RECOMMENDED FOR CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotherapy)

### NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying drugs
- Growth, gestation/lactation



#### STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



#### URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



#### LOW RSS

Helps lower the concentration of ions contributing to crystal formation.

**[S/O®] INDEX** This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	83.68 g	8 g
Fat content	62.76 g	6 g
Carbohydrate	48.12 g	4.6 g
Crude fibre	14.64 g	1.4 g
Dietary fibre	25.1 g	2.4 g
Calcium	2.2 g	0.21 g
Phosphorus	2.09 g	0.2 g
Potassium	2.09 g	0.2 g
Sodium	3.03 g	0.29 g
Chloride	4.18 g	0.4 g
Magnesium	0.17 g	0.02 g
Copper	3.14 mg	0.3 mg
Zinc	23.01 mg	2.2 mg
Taurine	1.26 g	0.12 g
Vitamin A	47071.13 IU	4500 IU
Vitamin E	198.74 mg	19 mg
Omega 6	25.1 g	2.4 g
Omega 3	1.36 g	0.13 g
EPA + DHA	0.63 g	0.06 g
Metabolisable energy*		96 kcal / 100 g

\*Calculated according to NRC2006.

### COMPOSITION

Meat and animal derivatives, oils and fats, cereals, derivatives of vegetable origin, minerals, various sugars.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 190 IU, E1 (Iron): 3.4 mg, E2 (Iodine): 0.24 mg, E4 (Copper): 2 mg, E5 (Manganese): 1 mg, E6 (Zinc): 10 mg, Urine acidifying substances: calcium sulfate: 0.25%.

Technological additives: Sodium bisulfate: 0.56 g.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



## URINARY S/O





## URINARY S/O



THIN SLICES IN GRAVY

Available in box of 12 pouches of 100 g

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

	DOG'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	pouch	grams	pouch	grams	pouch
ADULT DOG	2	207	2	182	2	157	1 + 1/2
	4	348	3 + 1/2	306	3	264	2 + 1/2
	6	471	4 + 1/2	415	4	358	3 + 1/2
	8	585	6	515	5	444	4 + 1/2
	10	691	7	608	6	525	5 + 1/2
	15	937	9 + 1/2	824	8	712	7
	20	1162	11 + 1/2	1023	10	883	9
	25	1374	13 + 1/2	1209	12	1044	10 + 1/2
	30	1576	16	1386	14	1197	12
	35	1769	17 + 1/2	1556	15 + 1/2	1344	13 + 1/2
	40	1955	19 + 1/2	1720	17	1486	15

Urinary S/O is a complete dietetic feed for dogs, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties, its low level of magnesium and its restricted level of protein but of high quality.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. Then the dog should be re-evaluated regularly by the veterinarian to assess the relevance of a life long recommendation. Water should be available at all times.

**NOTE:** -renal function should be checked before the advise of a new diet.

-in older dogs, the use of Urinary S/O Ageing<sup>7+</sup> is recommended.

## LOWER URINARY TRACT DISEASE



### RECOMMENDED FOR CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotherapy)

### NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying drugs
- Growth, gestation/lactation



#### STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



#### URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



#### LOW RSS

Helps lower the concentration of ions contributing to crystal formation.

[S/O<sup>®</sup> INDEX] This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	86.11 g	8 g
Fat content	59.2 g	5.5 g
Carbohydrate	51.67 g	4.8 g
Crude fibre	12.92 g	1.2 g
Dietary fibre	15.07 g	1.4 g
Calcium	1.61 g	0.15 g
Phosphorus	1.51 g	0.14 g
Potassium	1.61 g	0.15 g
Sodium	2.91 g	0.27 g
Chloride	3.98 g	0.37 g
Magnesium	0.13 g	0.01 g
Copper	3.77 mg	0.35 mg
Zinc	23.14 mg	2.15 mg
Taurine	1.24 g	0.12 g
Vitamin A	46286.33 IU	4300 IU
Vitamin E	226.05 mg	21 mg
Omega 6	19.38 g	1.8 g
Omega 3	1.35 g	0.13 g
EPA + DHA	0,65	0.06 g
Metabolisable energy*		93 kcal / 100g

\*Calculated according to NRC2006.

### COMPOSITION

Meat and animal derivatives, cereals, oils and fats, vegetable protein extracts, derivatives of vegetable origin, minerals, various sugars.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 210 IU, E1 (Iron): 3.5 mg, E2 (Iodine): 0.25 mg, E4 (Copper): 2 mg, E5 (Manganese): 1.1 mg, E6 (Zinc): 11 mg, Urine acidifying substances: calcium sulfate: 0.25%.

Technological additives: (acidity regulator) Sodium bisulfate: 1.107 g.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

## URINARY S/O



## URINARY S/O AGEING<sup>7+</sup>



Available in 1.5 kg, 3.5 kg, 8 kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

	DOG'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	cups	grams	cups	grams	cups
ADULT DOG	2	55	5/8	48	4/8	42	4/8
	4	92	1	81	7/8	70	6/8
	6	125	1 + 3/8	110	1 + 2/8	95	1
	8	155	1 + 6/8	136	1 + 4/8	117	1 + 2/8
	10	183	2	161	1 + 6/8	139	1 + 4/8
	15	248	2 + 6/8	218	2 + 3/8	188	2
	20	307	3 + 3/8	270	3	233	2 + 4/8
	25	363	4	320	3 + 4/8	276	3
	30	416	4 + 4/8	366	4	316	3 + 4/8
	35	467	5 + 1/8	411	4 + 4/8	355	3 + 7/8
	40	517	5 + 5/8	455	5	393	4 + 2/8
	45	564	6 + 1/8	497	5 + 4/8	429	4 + 6/8
	50	611	6 + 6/8	538	5 + 7/8	464	5 + 1/8
	55	656	7 + 2/8	577	6 + 3/8	499	5 + 4/8
	60	700	7 + 5/8	616	6 + 6/8	532	5 + 7/8
	70	786	8 + 5/8	692	7 + 5/8	597	6 + 4/8
	80	869	9 + 4/8	765	8 + 3/8	660	7 + 2/8

Urinary S/O Ageing<sup>7+</sup> is a complete dietetic feed for dogs, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties, its low level of magnesium and its restricted level of protein but of high quality.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O Ageing<sup>7+</sup> for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. Then the dog should be re-evaluated regularly by the veterinarian to assess the relevance of a life long recommendation. Water should be available at all times.

## LOWER URINARY TRACT DISEASE



### RECOMMENDED FOR AGEING DOGS (> 7 YEARS OLD) IN CASE OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotherapy)

### NOT RECOMMENDED IN CASE OF

- Hyperkalemia
- Concurrent use of urine-acidifying drugs
- Growth, gestation/lactation



#### URINARY TRACT HEALTH

Effective struvite stone dissolution & specifically formulated for up to 80% of all urinary stones.



#### AGEING SUPPORT

Support dogs' vitality and cognition with an exclusive blend of nutrients & antioxidants. Also support kidneys health.

[S/O<sup>®</sup> INDEX] This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	52 g	20 g
Fat content	44.18 g	17 g
Carbohydrate	114.35 g	44 g
Crude fibre	6.24 g	2.4 g
Dietary fibre	14.81 g	5.7 g
Calcium	1.74 g	0.67 g
Phosphorus	1.46 g	0.56 g
Potassium	4.55 g	1.75 g
Sodium	1.04 g	0.4 g
Chloride	3.82 g	1.47 g
Magnesium	0.23 g	0.09 g
Copper	3.9 mg	1.5 mg
Zinc	44.18 mg	17 mg
Taurine	0.49 g	0.19 g
Vitamin A	4417.88 IU	1700 IU
Vitamin E	124.74 mg	48 mg
Omega 6	8.97 g	3.45 g
Omega 3	2.39 g	0.92 g
EPA + DHA	1.04 g	0.4 g
Metabolisable energy*		385 kcal / 100g

\*Calculated according to NRC2006.

### COMPOSITION

Rice, maize flour, dehydrated poultry protein, animal fats, wheat flour, minerals, hydrolysed animal proteins, maize gluten, cellulose powder, wheat gluten\*\*, egg powder, fish oil, soya oil, dried tomato pulp (source of lycopene), fructo-oligo-saccharides, hydrolysed crustaceans (source of glucosamine), marigold extract (source of lutein), hydrolysed cartilage (source of chondroitin). Protein sources: dehydrated poultry protein, hydrolysed animal proteins, maize gluten, wheat gluten, egg powder.

\*\*L.I.P.: protein selected for its very high digestibility.

### ADDITIVES (KG)\*\*\*

Nutritional additives: Vitamin A: 15500 IU, Vitamin D3: 1000 IU, E1 (Iron): 60 mg, E2 (Iodine): 5.3 mg, E4 (Copper): 14 mg, E5 (Manganese): 67 mg, E6 (Zinc): 145 mg, E8 (Selenium): 0.11 mg, Urine acidifying substances: calcium sulphate (1.4%).

Technological additive: sodium bisulphate (urine acidifying substance): 5 g - Preservatives - Antioxidants.

\*\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

## URINARY S/O AGEING<sup>7+</sup>



## URINARY S/O AGEING 7+



Available in box of 12 pouches of 85 g

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

	DOG'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	pouch	grams	pouch	grams	pouch
ADULT DOG	2	203	2 + 1/2	178	2	154	2
	4	341	4	300	3 + 1/2	259	3
	6	462	5 + 1/2	406	5	351	4
	8	573	6 + 1/2	504	6	435	5
	10	677	8	596	7	515	6
	15	918	11	808	9 + 1/2	698	8
	20	1139	13 + 1/2	1002	12	866	10
	25	1346	16	1185	14	1023	12
	30	1544	18	1358	16	1173	14
	35	1733	20 + 1/2	1525	18	1317	15 + 1/2
	40	1915	22 + 1/2	1686	20	1456	17

Urinary S/O Ageing<sup>7+</sup> is a complete dietetic feed for dogs, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties, its low level of magnesium and its restricted level of protein but of high quality.

RECOMMENDATIONS: It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O Ageing<sup>7+</sup> for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. Then the dog should be re-evaluated regularly by the veterinarian to assess the relevance of a life long recommendation. Water should be available at all times.

## LOWER URINARY TRACT DISEASE



### RECOMMENDED FOR AGEING DOGS (> 7 YEARS OLD) IN CASE OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotherapy)

### NOT RECOMMENDED IN CASE OF

- Hyperkalemia
- Concurrent use of urine-acidifying drugs
- Growth, gestation/lactation



#### URINARY TRACT HEALTH

Effective struvite stone dissolution & specifically formulated for up to 80% of all urinary stones.



#### AGEING SUPPORT

Support dogs' vitality and cognition with an exclusive blend of nutrients & antioxidants. Also support kidneys health.



**[S/O® INDEX]** This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	91.68 g	8.7 g
Fat content	62.17 g	5.9 g
Carbohydrate	44.26 g	4.2 g
Crude fibre	17.91 g	1.7 g
Dietary fibre	23.18 g	2.2 g
Calcium	2 g	0.19 g
Phosphorus	1.48 g	0.14 g
Potassium	4.43 g	0.42 g
Sodium	1.05 g	0.1 g
Chloride	4.21 g	0.4 g
Magnesium	0.14 g	0.01 g
Copper	3.16 mg	0.3 mg
Zinc	27.4 mg	2.6 mg
Taurine	1.29 g	0.12 g
Vitamin A	50579.56 IU	4800 IU
Vitamin E	221.29 mg	21 mg
Omega 6	20.02 g	1.9 g
Omega 3	1.9 g	0.18 g
EPA + DHA	1.05 g	0.1 g
Metabolisable energy*		95 kcal / 100g

\*Calculated according to NRC2006.

### COMPOSITION

Meat and animal derivatives, derivatives of vegetable origin, cereals, oils and fats, minerals, various sugars, vegetables.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 210 IU, E1 (Iron): 4 mg, E2 (Iodine): 0.2 mg, E4 (Copper): 1.5 mg, E5 (Manganese): 1.3 mg, E6 (Zinc): 13 mg, Urine acidifying substances: calcium sulfate: 0.28%.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

## URINARY S/O

### MODERATE CALORIE



Available in 1.5 kg, 6.5 kg, 12.5 kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

	DOG'S WEIGHT (kg)	BODY CONDITION			
		NORMAL		OVERWEIGHT	
		grams	cups	grams	cups
ADULT DOG	2	49	5/8	41	4/8
	4	82	1	69	7/8
	6	111	1 + 3/8	93	1 + 1/8
	8	137	1 + 6/8	116	1 + 4/8
	10	163	2	137	1 + 6/8
	15	220	2 + 6/8	186	2 + 3/8
	20	273	3 + 4/8	230	2 + 7/8
	25	323	4 + 1/8	272	3 + 3/8
	30	370	4 + 5/8	312	4
	35	416	5 + 2/8	350	4 + 3/8
	40	460	5 + 6/8	387	4 + 7/8
	45	502	6 + 3/8	423	5 + 3/8
	50	543	6 + 7/8	458	5 + 6/8
	55	584	7 + 3/8	492	6 + 2/8
	60	623	7 + 7/8	525	6 + 5/8
	65	662	8 + 3/8	557	7
	70	699	8 + 7/8	589	7 + 3/8

Urinary S/O Moderate Calorie is a complete dietetic feed for dogs, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties, its low level of magnesium and its restricted level of protein but of high quality.

RECOMMENDATIONS: It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O Moderate Calorie for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. Then the dog should be re-evaluated regularly by the veterinarian to assess the relevance of a life long recommendation. Water should be available at all times.

NOTE: in older dogs, renal function should be checked before Urinary S/O Moderate Calorie is advised. It is recommended to divide the daily intake into 2 meals.

## LOWER URINARY TRACT DISEASE



### RECOMMENDED FOR DOGS WITH TENDENCY TO BE OVERWEIGHT IN CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotherapy)

### NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying drugs
- Growth, gestation/lactation



#### STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



#### MODERATE CALORIE

Moderate calorie content to help maintain ideal weight.



#### URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



#### LOW RSS

Helps lower the concentration of ions contributing to struvite and calcium oxalate crystal formation.



**[S/O® INDEX]** This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	60.85 g	20 g
Fat content	33.47 g	11 g
Carbohydrate	139.34 g	45.8 g
Crude fibre	19.77 g	6.5 g
Dietary fibre	37.12 g	12.2 g
Calcium	1.73 g	0.57 g
Phosphorus	1.73 g	0.57 g
Potassium	2.43 g	0.8 g
Sodium	3.65 g	1.2 g
Chloride	6.54 g	2.15 g
Magnesium	0.18 g	0.06 g
Copper	4.56 mg	1.5 mg
Zinc	51.72 mg	17 mg
Taurine	0.52 g	0.17 g
Vitamin A	6997.26 IU	2300 IU
Vitamin E	136.9 mg	45 mg
Omega 6	7.18 g	2.36g
Omega 3	1.52 g	0.5 g
EPA + DHA	0.61 g	0.2 g
Metabolisable energy*		329 kcal / 100g

\*Calculated according to NRC2006.

### COMPOSITION

Maize flour, rice, dehydrated poultry protein, vegetable fibres, maize gluten, animal fats, minerals, hydrolysed animal proteins, chicory pulp, wheat gluten\*\*, soya oil, fish oil, fructo-oligo-saccharides, mono - and diglycerides of palmitic and stearic acids esterified with citric acid, marigold extract (source of lutein). Protein sources: dehydrated poultry protein, maize gluten, hydrolysed animal proteins, wheat gluten.

\*\*L.I.P.: protein selected for its very high digestibility

### ADDITIVES (KG)\*\*\*

Nutritional additives: Vitamin A: 21500 IU, Vitamin D3: 1400 IU, E1 (Iron): 44 mg, E2 (Iodine): 4.4 mg, E4 (Copper): 14 mg, E5 (Manganese): 58 mg, E6 (Zinc): 146 mg, E8 (Selenium): 0.11 mg, Urine acidifying substances: calcium sulphate (1%). Preservatives - Antioxidants.

\*\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

URINARY S/O AGEING 7+



URINARY S/O MODERATE CALORIE





## URINARY S/O

MODERATE CALORIE



THIN SLICES IN GRAVY

Available in box of 12 pouches of 100 g

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

	DOG'S WEIGHT (kg)	BODY CONDITION			
		NORMAL		OVERWEIGHT	
		grams	pouch	grams	pouch
ADULT DOG	2	181	2	152	1 + 1/2
	4	304	3	256	2 + 1/2
	6	412	4	347	3 + 1/2
	8	512	5	431	4 + 1/2
	10	605	6	509	5
	15	820	8	691	7
	20	1018	10	857	8 + 1/2
	25	1203	12	1013	10
	30	1379	14	1161	11 + 1/2
	35	1548	15 + 1/2	1304	13
	40	1711	17	1441	14 + 1/2

Urinary S/O Moderate Calorie is a complete dietetic feed for dogs, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties, its low level of magnesium and its restricted level of protein but of high quality.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O Moderate Calorie for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. Then the dog should be re-evaluated regularly by the veterinarian to assess the relevance of a life long recommendation. Water should be available at all times.

**IN OLDER DOGS:** renal function should be checked before Urinary S/O Moderate Calorie is advised. It is recommended to divide the daily intake into 2 meals.

## LOWER URINARY TRACT DISEASE



### RECOMMENDED FOR DOGS WITH TENDENCY TO BE OVERWEIGHT IN CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotherapy)

### NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying drugs
- Growth, gestation/lactation



#### STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



#### MODERATE CALORIE

Moderate calorie content to help maintain ideal weight.



#### URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



#### LOW RSS

Helps lower the concentration of ions contributing to crystal formation.



**[S/O®] INDEX** This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	97.32 g	8 g
Fat content	43.8 g	3.6 g
Carbohydrate	71.78 g	5.9 g
Crude fibre	12.17 g	1 g
Dietary fibre	17.64 g	1.5 g
Calcium	1.7 g	0.14 g
Phosphorus	1.58 g	0.13 g
Potassium	1.82 g	0.15 g
Sodium	2.92 g	0.24 g
Chloride	4.26 g	0.35 g
Magnesium	0.12 g	0.01 g
Copper	3.65 mg	0.30 mg
Zinc	21.29 mg	1.75 mg
Taurine	1.22 g	0.1 g
Vitamin A	48661.8 IU	4000 IU
Vitamin E	182.48 mg	15 mg
Omega 6	973 g	0.80 g
Omega 3	1.34 g	0.11 g
EPA + DHA	0.61 g	0.05 g
Metabolisable energy*		82 kcal / 100 g

\*Calculated according to NRC2006.

### COMPOSITION

Meat and animal derivatives, cereals, vegetable protein extracts, derivatives of vegetable origin, minerals, oils and fats, various sugars.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 180 IU, E1 (Iron): 2 mg, E2 (Iodine): 0.2 mg, E4 (Copper): 1.8 mg, E5 (Manganese): 0.8 mg, E6 (Zinc): 8 mg, Sulfur: 0.2%, Urine acidifying substances: calcium sulfate: 0.3%. Technological additives: (acidity regulator) Sodium bisulfate: 1.1 g.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

## URINARY S/O

SMALL DOGS



Available in 1.5 kg, 4 kg, 8 kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

	DOG'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	cups	grams	cups	grams	cups
ADULT DOG	2	55	5/8	48	4/8	42	4/8
	2,5	65	5/8	57	5/8	49	4/8
	3	74	6/8	65	5/8	56	5/8
	3,5	83	7/8	73	6/8	63	5/8
	4	92	1	81	7/8	70	6/8
	5	109	1 + 1/8	96	1	83	7/8
	6	125	1 + 2/8	110	1 + 1/8	95	1
	7	140	1 + 4/8	123	1 + 2/8	106	1 + 1/8
	8	155	1 + 5/8	136	1 + 3/8	117	1 + 2/8
	9	169	1 + 6/8	149	1 + 4/8	128	1 + 3/8
	10	183	1 + 7/8	161	1 + 5/8	139	1 + 4/8

Urinary S/O Small Dog is a complete dietetic feed for dogs, formulated to dissolve struvite stones and reducing their recurrence through its urine acidifying properties, its low level of magnesium and its restricted level of protein but of high quality.

**RECOMMENDATIONS:** It is recommended that a veterinarian's opinion be sought before use. Feed Urinary S/O Small Dog for 5 to 12 weeks for the dissolution of struvite stones and up to 6 months for the reduction of struvite stone recurrence. Then the dog should be re-evaluated regularly by the veterinarian to assess the relevance of a life long recommendation. Water should be available at all times.

**NOTE:** - renal function should be checked before the advice of a new diet.  
- in older dogs, the use of Urinary S/O Ageing® is recommended.

## LOWER URINARY TRACT DISEASE



### RECOMMENDED FOR CASES OF

- Struvite uroliths: dissolution and management of recurrence
- Calcium oxalate uroliths: management of recurrence
- Bacterial cystitis: management of secondary struvite crystalluria (in association with adequate antibiotherapy)

### NOT RECOMMENDED IN CASE OF

- Chronic Kidney Disease (CKD)
- Heart disease (when sodium restriction is sought)
- Concurrent use of urine-acidifying drugs
- Growth, gestation/lactation



#### STRUVITE DISSOLUTION

Helps dissolve all types of struvite stones.



#### LOW RSS

Helps lower the concentration of ions contributing to crystal formation.



#### URINE DILUTION

Urine dilution makes the urine less liable to form struvite and calcium oxalate stones.



#### TARTAR CONTROL

Nutrient that traps the calcium in saliva so reducing tartar deposits.



**[S/O®] INDEX** This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	51.98 g	20 g
Fat content	44.18 g	17 g
Carbohydrate	114.09 g	43.9 g
Crude fibre	5.98 g	2.3 g
Dietary fibre	14.55 g	5.6 g
Calcium	1.85 g	0.71 g
Phosphorus	1.82 g	0.7 g
Potassium	2.08 g	0.8 g
Sodium	3.12 g	1.2 g
Chloride	5.2 g	2 g
Magnesium	0.13 g	0.05 g
Copper	3.9 mg	1.5 mg
Zinc	44.18 mg	17 mg
Taurine	0.49 g	0.19 g
Vitamin A	4547.82 IU	1750 IU
Vitamin E	124.74 mg	48 mg
Omega 6	9.3 g	3.58 g
Omega 3	1.72 g	0.66 g
EPA + DHA	0.57 g	0.22 g
Metabolisable energy*		385 kcal / 100 g

\*Calculated according to NRC2006.

### COMPOSITION

Rice, maize flour, dehydrated poultry protein, animal fats, maize gluten, hydrolysed animal proteins, minerals, vegetable fibres, egg powder, wheat gluten\*\*, soya oil, fish oil, fructo-oligo-saccharides, marigold extract (source of lutein). Protein sources : dehydrated poultry protein, maize gluten, hydrolysed animal proteins, egg powder.

\*\*L.I.P.: protein selected for its very high digestibility.

### ADDITIVES (KG)\*\*\*

Nutritional additives: Vitamin A: 16000 IU, Vitamin D3: 1000 IU, E1 (Iron): 47 mg, E2 (Iodine): 4.7 mg, E4 (Copper): 14 mg, E5 (Manganese): 61 mg, E6 (Zinc): 148 mg, E8 (Selenium): 0.06 mg, Urine acidifying substance: calcium sulphate (0.8%). Technological additives: Pentasodium triphosphate: 3.5 g. Preservatives - Antioxidants.

\*\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

URINARY S/O MODERATE CALORIE



S

SMALL DOGS

URINARY S/O



URINARY U/C



Available in 2 kg, 7.5 kg, 14 kg bags

RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	cups	grams	cups	grams	cups
	2	55	5/8	48	4/8	42	4/8
	4	92	1	81	7/8	70	6/8
	6	125	1 + 3/8	110	1 + 2/8	95	1
	8	155	1 + 6/8	137	1 + 4/8	118	1 + 2/8
	10	184	2	162	1 + 6/8	140	1 + 4/8
	15	249	2 + 6/8	219	2 + 3/8	189	2 + 1/8
	20	309	3 + 3/8	272	3	235	2 + 5/8
	25	365	4	321	3 + 4/8	278	3
	30	419	4 + 5/8	368	4	318	3 + 4/8
	35	470	5 + 1/8	414	4 + 4/8	357	3 + 7/8
	40	520	5 + 6/8	457	5	395	4 + 3/8
	45	568	6 + 2/8	499	5 + 4/8	431	4 + 6/8
	50	614	6 + 6/8	541	5 + 7/8	467	5 + 1/8
	55	660	7 + 2/8	581	6 + 3/8	501	5 + 4/8
	60	704	7 + 6/8	620	6 + 6/8	535	5 + 7/8
	70	791	8 + 5/8	696	7 + 5/8	601	6 + 5/8
	80	874	9 + 5/8	769	8 + 3/8	664	7 + 2/8

Urinary U/C is a complete dietetic feed for dogs, formulated to reduce the formation of urate and cystine stones through its low level of protein and of purines, its high quality protein, its moderate level of sulphur amino acids and its urine alkalinising properties.

RECOMMENDATIONS: Feed Urinary U/C for up to 6 months to reduce the formation of urate stones, for up to 1 year to reduce the formation of cystine stones and for life in the case of irreversible disturbance of uric acid metabolism. It is recommended that a veterinarian's opinion be sought before use or before extending the period of use. Then the dog should be re-evaluated regularly by the veterinarian to assess the relevance of a life long recommendation. Water should be available at all times.

LOWER URINARY TRACT DISEASE



RECOMMENDED FOR CASES OF

- Metabolic uroliths (urate, xanthine, cystine): management of recurrence
  - Hyperuricosuria
  - Management of metabolic uroliths risk in dogs with leishmaniasis treated with xanthine oxidase inhibitors (eg allopurinol)\*
- \*If they present a secondary kidney disease (protein losing nephropathy with or without azotemia) then switch to a Renal diet*

NOT RECOMMENDED IN CASE OF

- Growth, gestation/lactation



URATE CONTROL

A selection of proteins with low purine content helps reduce the formation of urate urinary stones.



CYSTINE CONTROL

A limited intake of proteins and of some amino acids helps limit the formation of cystine urinary stones.



SKIN BARRIER

A complex to support the barrier effect of the skin.



ANTIOXIDANT COMPLEX

A synergistic complex of antioxidants to help neutralise free radicals.

KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	47.03 g	18 g
Fat content	39.2 g	15 g
Carbohydrate	130.65 g	50 g
Crude fibre	5.75 g	2.2 g
Dietary fibre	14.89 g	5.7 g
Calcium	2.06 g	0.79 g
Phosphorus	1.54 g	0.59 g
Potassium	2.35 g	0.9 g
Sodium	0.78 g	0.3 g
Chloride	1.44 g	0.55 g
Magnesium	0.1 g	0.04 g
Copper	3.92 mg	1.5 mg
Zinc	44.42 mg	17 mg
Taurine	0.55 g	0.21 g
Vitamin A	7839.04 IU	3000 IU
Vitamin E	159.39 mg	61 mg
Omega 6	7.24 g	2.77 g
Omega 3	1.59 g	0.61 g
EPA + DHA	0.68 g	0.26 g
Metabolisable energy*		383 kcal / 100 g

\*Calculated according to NRC2006.

COMPOSITION

Rice, maize flour, wheat flour, animal fats, egg powder, wheat gluten\*\*, maize gluten, minerals, vegetable fibres, fish oil, soya oil, fructo-oligo-saccharides, borage oil, marigold extract (source of lutein). Protein sources: egg powder, wheat gluten, maize gluten.

\*\*L.I.P.: protein selected for its very high digestibility.

ADDITIVES (KG)\*\*\*

Nutritional additives: Vitamin A: 30000 IU, Vitamin D3: 800 IU, E1 (Iron): 63 mg, E2 (Iodine): 5.3 mg, E4 (Copper): 14 mg, E5 (Manganese): 66 mg, E6 (Zinc): 147 mg, E8 (Selenium): 0.16 mg - Preservatives - Antioxidants.

\*\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

URINARY U/C



# DOGS MIX FEEDING GUIDELINES



**MIX: URINARY S/O DRY +  
URINARY S/O LOAF 400G WET**

DOG'S WEIGHT (kg)	CAN	WEIGHT CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)
2	1/4+	28	3/8 cup	21	2/8 cup	15	1/8 cup
4	1/4+	64	6/8 cup	54	5/8 cup	43	4/8 cup
6	1/2+	70	6/8 cup	55	5/8 cup	41	4/8 cup
8	1/2+	100	1 cup + 1/8 cup	82	1 cup	63	6/8 cup
10	1/2+	128	1 cup + 4/8 cup	106	1 cup + 2/8 cup	84	1 cup
15	1+	139	1 cup + 5/8 cup	109	1 cup + 2/8 cup	80	7/8 cup
20	1+	198	2 cup + 2/8 cup	161	1 cup + 7/8 cup	125	1 cup + 4/8 cup
25	1+	254	2 cup + 7/8 cup	210	2 cup + 3/8 cup	167	1 cup + 7/8 cup
30	1+	306	3 cup + 4/8 cup	257	3 cup	207	2 cup + 3/8 cup
40	1+	406	4 cup + 6/8 cup	344	4 cup	283	3 cup + 2/8 cup
50	1+	499	5 cup + 6/8 cup	426	4 cup + 7/8 cup	354	4 cup + 1/8 cup
60	1+	588	6 cup + 6/8 cup	504	5 cup + 7/8 cup	421	4 cup + 7/8 cup
70	1+	673	7 cup + 6/8 cup	579	6 cup + 6/8 cup	486	5 cup + 5/8 cup
80	1+	755	8 cup + 6/8 cup	652	7 cup + 4/8 cup	548	6 cup + 3/8 cup



**MIX: URINARY S/O DRY +  
URINARY S/O LOAF 200G WET**

DOG'S WEIGHT (kg)	CAN	WEIGHT CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)
2	1/2+	28	3/8 cup	22	2/8 cup	15	1/8 cup
4	1/2+	65	6/8 cup	54	5/8 cup	43	4/8 cup
6	1+	71	7/8 cup	57	5/8 cup	42	4/8 cup
8	1+	101	1 cup + 1/8 cup	83	1 cup	64	6/8 cup
10	1+	129	1 cup + 4/8 cup	107	1 cup + 2/8 cup	86	1 cup
15	1+	193	2 cup + 2/8 cup	164	1 cup + 7/8 cup	135	1 cup + 4/8 cup
20	1+	253	2 cup + 7/8 cup	216	2 cup + 4/8 cup	180	2 cup + 1/8 cup
25	1+	308	3 cup + 5/8 cup	265	3 cup + 1/8 cup	222	2 cup + 5/8 cup
30	1+	361	4 cup + 1/8 cup	311	3 cup + 5/8 cup	262	3 cup
40	1+	460	5 cup + 3/8 cup	399	4 cup + 5/8 cup	337	3 cup + 7/8 cup
50	1+	554	6 cup + 3/8 cup	481	5 cup + 5/8 cup	408	4 cup + 6/8 cup
60	1+	642	7 cup + 3/8 cup	559	6 cup + 4/8 cup	476	5 cup + 4/8 cup
70	1+	727	8 cup + 3/8 cup	634	7 cup + 3/8 cup	540	6 cup + 2/8 cup
80	1+	810	9 cup + 3/8 cup	706	8 cup + 1/8 cup	603	7 cup

# DOGS MIX FEEDING GUIDELINES



**MIX: URINARY S/O DRY +  
URINARY S/O CIG WET**

DOG'S WEIGHT (kg)	POUCH	WEIGHT CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)
2	1+	28	3/8 cup	21	2/8 cup	15	1/8 cup
4	1+	65	6/8 cup	54	5/8 cup	43	4/8 cup
6	1+	97	1 cup + 1/8 cup	82	1 cup	68	6/8 cup
8	1+	127	1 cup + 4/8 cup	109	1 cup + 2/8 cup	90	1 cup
10	1+	155	1 cup + 6/8 cup	133	1 cup + 4/8 cup	111	1 cup + 2/8 cup
15	1+	219	2 cup + 4/8 cup	190	2 cup + 2/8 cup	160	1 cup + 7/8 cup
20	1+	278	3 cup + 2/8 cup	242	2 cup + 6/8 cup	205	2 cup + 3/8 cup
25	1+	334	3 cup + 7/8 cup	291	3 cup + 3/8 cup	247	2 cup + 7/8 cup
30	1+	387	4 cup + 4/8 cup	337	3 cup + 7/8 cup	288	3 cup + 3/8 cup
40	1+	486	5 cup + 5/8 cup	425	4 cup + 7/8 cup	363	4 cup + 2/8 cup
50	1+	579	6 cup + 6/8 cup	507	5 cup + 7/8 cup	434	5 cup
60	1+	668	7 cup + 6/8 cup	585	6 cup + 6/8 cup	502	5 cup + 6/8 cup
70	1+	753	8 cup + 6/8 cup	660	7 cup + 5/8 cup	566	6 cup + 4/8 cup
80	1+	835	9 cup + 5/8 cup	732	8 cup + 4/8 cup	629	7 cup + 2/8 cup



**MIX: URINARY S/O AGEING 7+ DRY +  
URINARY S/O AGEING 7+ LOAF WET**

DOG'S WEIGHT (kg)	POUCH	WEIGHT CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)
2	1+	32	3/8 cup	25	2/8 cup	19	2/8 cup
4	1+	69	6/8 cup	58	5/8 cup	47	4/8 cup
6	1+	102	1 cup + 1/8 cup	87	1 cup	72	6/8 cup
8	1+	132	1 cup + 4/8 cup	113	1 cup + 2/8 cup	95	1 cup
10	1+	160	1 cup + 6/8 cup	138	1 cup + 4/8 cup	116	1 cup + 2/8 cup
15	1+	225	2 cup + 4/8 cup	195	2 cup + 1/8 cup	165	1 cup + 6/8 cup
20	1+	284	3 cup + 1/8 cup	247	2 cup + 6/8 cup	211	2 cup + 2/8 cup
25	1+	340	3 cup + 6/8 cup	297	3 cup + 2/8 cup	253	2 cup + 6/8 cup
30	1+	393	4 cup + 3/8 cup	344	3 cup + 6/8 cup	294	3 cup + 2/8 cup
40	1+	494	5 cup + 3/8 cup	432	4 cup + 6/8 cup	370	4 cup
50	1+	588	6 cup + 4/8 cup	515	5 cup + 5/8 cup	441	4 cup + 7/8 cup
60	1+	677	7 cup + 3/8 cup	593	6 cup + 4/8 cup	509	5 cup + 5/8 cup
70	1+	763	8 cup + 3/8 cup	669	7 cup + 3/8 cup	575	6 cup + 2/8 cup
80	1+	846	9 cup + 2/8 cup	742	8 cup + 1/8 cup	637	7 cup





# DOGS MIX FEEDING GUIDELINES



## MIX: URINARY S/O MODERATE CALORIE DRY + URINARY S/O MODERATE CALORIE CIG WET

DOG'S WEIGHT (kg)	POUCH	WEIGHT CONDITION			
		NORMAL		OVERWEIGHT	
		DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)
2	1+	22	2/8 cup	14	1/8 cup
4	1+	55	6/8 cup	42	4/8 cup
6	1+	84	1 cup	66	7/8 cup
8	1+	111	1 cup + 3/8 cup	89	1 cup + 1/8 cup
10	1+	136	1 cup + 6/8 cup	110	1 cup + 3/8 cup
15	1+	193	2 cup + 4/8 cup	159	2 cup
20	1+	246	3 cup + 1/8 cup	203	2 cup + 5/8 cup
25	1+	296	3 cup + 6/8 cup	245	3 cup + 1/8 cup
30	1+	344	4 cup + 3/8 cup	285	3 cup + 5/8 cup
40	1+	433	5 cup + 4/8 cup	360	4 cup + 4/8 cup
50	1+	517	6 cup + 4/8 cup	431	5 cup + 4/8 cup
60	1+	596	7 cup + 4/8 cup	498	6 cup + 2/8 cup
70	1+	673	8 cup + 4/8 cup	562	7 cup + 1/8 cup
80	1+	746	9 cup + 3/8 cup	624	7 cup + 7/8 cup




## MIX: URINARY S/O SMALL DOG DRY + URINARY S/O CIG WET

DOG'S WEIGHT (kg)	POUCH	WEIGHT CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)
2	1+	28	2/8 cup	22	2/8 cup	15	1/8 cup
4	1+	65	5/8 cup	54	5/8 cup	43	4/8 cup
6	1+	98	1 cup	83	7/8 cup	68	6/8 cup
8	1+	128	1 cup + 3/8 cup	110	1 cup + 1/8 cup	91	1 cup
10	1+	156	1 cup + 5/8 cup	134	1 cup + 3/8 cup	112	1 cup + 1/8 cup



## MIX: URINARY S/O SMALL DOG DRY + URINARY S/O LOAF 400G WET

DOG'S WEIGHT (kg)	CAN	WEIGHT CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)
2	1/4+	28	2/8 cup	21	2/8 cup	15	1/8 cup
4	1/4+	65	5/8 cup	54	4/8 cup	43	4/8 cup
6	1/2+	71	6/8 cup	56	5/8 cup	41	3/8 cup
8	1/2+	101	1 cup	82	7/8 cup	64	5/8 cup
10	1/2+	129	1 cup + 3/8 cup	107	1 cup + 1/8 cup	85	7/8 cup



## ANALLERGENIC



Available in 1.5 kg, 3 kg, 8 kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

DOG'S WEIGHT (kg)	BODY CONDITION					
	THIN		NORMAL		OVERWEIGHT	
	grams	cups	grams	cups	grams	cups
2	53	5/8	46	4/8	40	4/8
4	88	1	78	7/8	67	6/8
6	120	1 + 3/8	105	1 + 1/8	91	1
8	149	1 + 5/8	131	1 + 3/8	113	1 + 2/8
10	176	1 + 7/8	155	1 + 6/8	133	1 + 4/8
15	238	2 + 5/8	209	2 + 2/8	181	2
20	295	3 + 2/8	260	2 + 7/8	224	2 + 4/8
25	349	3 + 7/8	307	3 + 3/8	265	2 + 7/8
30	400	4 + 3/8	352	3 + 7/8	304	3 + 3/8
35	449	4 + 7/8	395	4 + 3/8	341	3 + 6/8
40	497	5 + 4/8	437	4 + 6/8	377	4 + 1/8
45	543	6	477	5 + 2/8	412	4 + 4/8
50	587	6 + 3/8	517	5 + 5/8	446	4 + 7/8
55	631	6 + 7/8	555	6 + 1/8	479	5 + 2/8
60	673	7 + 3/8	592	6 + 4/8	512	5 + 5/8
70	756	8 + 2/8	665	7 + 2/8	574	6 + 2/8
80	835	9 + 1/8	735	8	635	7

When a food allergy or a food intolerance is suspected, the dog's diet should be changed without any food transition. Nutritional management with Anallergenic may be continued for the pet's lifetime. It is recommended to divide the daily intake into 2 meals.

## ADVERSE REACTION TO FOOD



### RECOMMENDED FOR CASES OF

- Adverse Food Reactions (AFR) with dermatologic and/or gastro-intestinal signs:
  - diagnosis: food elimination trial
  - management
- Canine Atopic Dermatitis (CAD) associated with AFR
- Inflammatory Bowel Disease (IBD)

### NOT RECOMMENDED IN CASE OF

- Pancreatitis
- Growth, gestation/lactation



#### OLIGOPEPTIDES

Low molecular weight peptides to reduce the risk of allergic reactions to food.



#### ALLERGEN RESTRICTION

Formula and production process aimed at excluding sources of food allergens.



#### SKIN BARRIER

Formulated to support the skin's natural protective barrier for optimal skin health.



#### ANTIOXIDANT COMPLEX

A synergistic complex of antioxidants to help neutralise free radicals.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	45 g	18 g
Fat content	41.2 g	16.5 g
Carbohydrate	126.9 g	50.8 g
Crude fibre	4.2 g	1.7 g
Dietary fibre	12.5 g	5 g
Calcium	1.5 g	0.62 g
Phosphorus	1.2 g	0.48 g
Potassium	1.9 g	0.75 g
Sodium	1.7 g	0.7 g
Chloride	3.5 g	1.41 g
Magnesium	0.2 g	0.08 g
Copper	3.7 mg	1.5 mg
Zinc	44.2 mg	17.7 mg
Taurine	0.5 g	0.19 g
Vitamin A	9118.2 IU	3650 IU
Vitamin E	174.9 mg	70 mg
Omega 6	7.5 g	3.01 g
Omega 3	1.3 g	0.52 g
EPA + DHA	0.5 g	0.2 g
Metabolisable energy*		400 kcal/100 g

\*Calculated according to NRC2006.

### COMPOSITION

Maize starch, feather hydrolysate with low molecular weight (source of L amino acids and oligopeptides), copra oil, soya oil, minerals, vegetable fibres, chicory pulp, fructo-oligo-saccharides, fish oil, mono - and diglycerides of palmitic and stearic acids esterified with citric acid, animal fat, marigold extract (source of lutein). Protein source: feather hydrolysate with low molecular weight. Carbohydrate source: maize starch.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin A: 36500 IU, Vitamin D3: 800 IU, E1 (Iron): 42 mg, E2 (Iodine): 3.8 mg, E4 (Copper): 15 mg, E5 (Manganese): 55 mg, E6 (Zinc): 154 mg, E8 (Selenium): 0.26 mg - Technological additives: Clinoptilolite of sedimentary origin: 8.48 g - Preservatives - Antioxidants.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



## ANALLERGENIC



## HYPOALLERGENIC



Available in 2 kg, 7 kg, 14 kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

DOG'S WEIGHT (kg)	BODY CONDITION					
	THIN		NORMAL		OVERWEIGHT	
	grams	cups	grams	cups	grams	cups
2	52	5/8	45	4/8	39	4/8
4	87	1	76	7/8	66	6/8
6	117	1 + 3/8	103	1 + 2/8	89	1
8	146	1 + 6/8	128	1 + 4/8	111	1 + 2/8
10	172	2	152	1 + 6/8	131	1 + 4/8
15	234	2 + 6/8	206	2 + 3/8	178	2
20	290	3 + 3/8	255	3	220	2 + 4/8
25	343	4	302	3 + 4/8	260	3
30	393	4 + 4/8	346	4	299	3 + 4/8
35	441	5 + 1/8	388	4 + 4/8	335	3 + 7/8
40	487	5 + 5/8	429	5	370	4 + 2/8
45	532	6 + 1/8	469	5 + 3/8	405	4 + 6/8
50	576	6 + 5/8	507	5 + 7/8	438	5 + 1/8
55	619	7 + 1/8	545	6 + 2/8	470	5 + 4/8
60	661	7 + 5/8	581	6 + 6/8	502	5 + 6/8
70	742	8 + 5/8	653	7 + 4/8	564	6 + 4/8
80	820	9 + 4/8	721	8 + 3/8	623	7 + 2/8

When a food allergy or a food intolerance is suspected, the dog's diet could be changed without any food transition. Nutritional management with Hypoallergenic may be continued for the pet's lifetime. It is recommended to divide the daily intake into 2 meals.

NOTE: For the mixed feeding table please see the end of the section Dermatology.

## ADVERSE REACTION TO FOOD



### RECOMMENDED FOR CASES OF

- Adverse Food Reactions (AFR) with dermatologic and/or gastrointestinal signs:
  - diagnosis: food elimination trial
  - management
- Canine Atopic Dermatitis (CAD) + chronic pruritus
- Chronic diarrhea
- Inflammatory Bowel Disease (IBD)
- Exocrine Pancreatic Insufficiency (EPI)
- Small intestinal bacterial overgrowth (SIBO)

### NOT RECOMMENDED IN CASE OF

- Pancreatitis
- Gestation/lactation



#### HYDROLYSED PROTEIN

Hydrolysed protein with low molecular weight to ensure the food is hypoallergenic.



#### SKIN BARRIER

Formulated to support the skin's natural protective barrier for optimal skin health.



#### EPA/DHA

Fatty acids to help maintain a healthy digestive system and a healthy skin.



#### DIGESTIVE HEALTH

Nutrients which support a balanced digestive system.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	51.5 g	21 g
Fat content	46.6 g	19 g
Carbohydrate	106.2 g	43.3 g
Crude fibre	2.7 g	1.1 g
Dietary fibre	13.7 g	5.6 g
Calcium	2 g	0.8 g
Phosphorus	1.5 g	0.6 g
Potassium	1.5 g	0.6 g
Sodium	1 g	0.4 g
Chloride	1.9 g	0.76 g
Magnesium	0.2 g	0.09 g
Copper	3.7 mg	1.5 mg
Zinc	41.7 mg	17 mg
Taurine	0.5 g	0.21 g
Vitamin A	7232.2 IU	2950 IU
Vitamin E	149.5 mg	61 mg
Omega 6	10.9 g	4.46 g
Omega 3	2.3 g	0.95 g
EPA + DHA	0.8 g	0.33 g
Metabolisable energy*		408 kcal/100 g

\*Calculated according to NRC2006.

### COMPOSITION

Rice, hydrolysed soya protein isolate, animal fats, hydrolysed poultry liver, beet pulp, minerals, soya oil, fish oil, fructo-oligo-saccharides, borage oil, marigold extract (source of lutein). Protein sources: hydrolysed soya protein isolate, hydrolysed poultry liver. Carbohydrate source: rice.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin A: 27500 IU, Vitamin D3: 800 IU, E1 (Iron): 42 mg, E2 (Iodine): 3.8 mg, E4 (Copper): 15 mg, E5 (Manganese): 55 mg, E6 (Zinc): 154 mg, E8 (Selenium): 0.26 mg - Technological additives: Clinoptilolite of sedimentary origin: 10 g - Preservatives - Antioxidants.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



## HYPOALLERGENIC





## HYPOALLERGENIC



LOAF



Available in 200 g can

### RECOMMENDED DAILY INTAKE & FEEDING DURATION BASED ON 200 G

DOG'S WEIGHT (kg)	BODY CONDITION					
	THIN		NORMAL		OVERWEIGHT	
	grams	can	grams	can	grams	can
2	198	1	175	1	151	3/4
4	334	1 + 3/4	294	1 + 2/4	253	1 + 1/4
6	452	2 + 1/4	398	2	344	1 + 3/4
8	561	2 + 3/4	494	2 + 2/4	426	2 + 1/4
10	663	3 + 1/4	584	3	504	2 + 2/4
15	899	4 + 2/4	791	4	683	3 + 2/4
20	1 115	5 + 2/4	981	5	848	4 + 1/4
25	1 318	6 + 2/4	1 160	5 + 3/4	1 002	5
30	1 512	7 + 2/4	1 330	6 + 3/4	1 149	5 + 3/4
35	1 697	8 + 2/4	1 493	7 + 2/4	1 290	6 + 2/4
40	1 876	9 + 2/4	1 651	8 + 1/4	1 425	7 + 1/4
45	2 049	10 + 1/4	1 803	9	1 557	7 + 3/4
50	2 217	11	1 951	9 + 3/4	1 685	8 + 2/4
55	2 382	12	2 096	10 + 2/4	1 810	9
60	2 542	12 + 3/4	2 237	11 + 1/4	1 932	9 + 3/4
70	2 854	14 + 1/4	2 511	12 + 2/4	2 169	10 + 3/4
80	3 154	15 + 3/4	2 776	14	2 397	12

When a food allergy or a food intolerance is suspected, the dog's diet could be changed without any food transition. The hypoallergenic nutritional management may be continued for the pet's lifetime. It is recommended to divide the daily intake into 2 meals. For the mixed feeding table please see the end of the section "Adverse reaction to food".

NOTE: For the mixed feeding table please see the end of the section Dermatology.

## ADVERSE REACTION TO FOOD



### RECOMMENDED FOR CASES OF

- Adverse Food Reactions (AFR) with dermatologic and/or gastrointestinal signs:
  - diagnosis: food elimination trial
  - management
- Canine Atopic Dermatitis (CAD) associated with AFR
- Chronic diarrhea
- Inflammatory Bowel Disease (IBD)
- Exocrine Pancreatic Insufficiency (EPI)
- Small intestinal bacterial overgrowth (SIBO)

### NOT RECOMMENDED IN CASE OF

- Pancreatitis
- Growth, gestation/lactation



#### HYDROLYSED PROTEIN

Hydrolysed protein with low molecular weight to ensure the food is hypoallergenic.



#### SKIN BARRIER

Formulated to support the skin's natural protective barrier for optimal skin health.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	64.8 g	6.4 g
Fat content	35.4 g	3.5 g
Carbohydrate	127.5 g	12.6 g
Crude fibre	15.2 g	1.5 g
Dietary fibre	22.3 g	2.2 g
Calcium	2 g	0.2 g
Phosphorus	1.6 g	0.16 g
Potassium	2 g	0.2 g
Sodium	2.4 g	0.24 g
Chloride	2 g	0.2 g
Magnesium	0.2 g	0.02 g
Copper	5.1 mg	0.5 mg
Zinc	45.5 mg	4.5 mg
Taurine	1.2 g	0.12 g
Vitamin A	10121.5 IU	1000 IU
Vitamin E	151.18 mg	15 mg
Omega 6	16.2 g	1.6 g
Omega 3	1.7 g	0.17 g
EPA + DHA	1.2 g	0.12 g
Metabolisable energy*		99 kcal/100 g

\*Calculated according to NRC2006.

### COMPOSITION

Derivatives of vegetable origin, vegetable protein extracts, oils and fats, meat and animal derivatives, minerals. Selected sources of protein: hydrolysed soya protein concentrate, hydrolysed poultry by-products. Selected source of carbohydrate: pea starch.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin A: 3000 IU, Vitamin D3: 320 IU, E1 (Iron): 11 mg, E2 (Iodine): 1 mg, E4 (Copper): 3.8 mg, E5 (Manganese): 15 mg, E6 (Zinc): 36 mg - Technological additives: Clinoptilolite of sedimentary origin: 2.5 g.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

## HYPOALLERGENIC



## HYPOALLERGENIC



LOAF



Available in 400 g can

### RECOMMENDED DAILY INTAKE & FEEDING DURATION BASED ON 400 G

DOG'S WEIGHT (kg)	BODY CONDITION					
	THIN		NORMAL		OVERWEIGHT	
	grams	can	grams	can	grams	can
2	198	2/4	175	2/4	151	2/4
4	334	3/4	294	3/4	253	3/4
6	452	1 + 1/4	398	1	344	3/4
8	561	1 + 2/4	494	1 + 1/4	426	1
10	663	1 + 3/4	584	1 + 2/4	504	1 + 1/4
15	899	2 + 1/4	791	2	683	1 + 3/4
20	1115	2 + 3/4	981	2 + 2/4	848	2
25	1318	3 + 1/4	1160	3	1002	2 + 2/4
30	1512	3 + 3/4	1330	3 + 1/4	1149	2 + 3/4
35	1697	4 + 1/4	1493	3 + 3/4	1290	3 + 1/4
40	1876	4 + 3/4	1651	4 + 1/4	1425	3 + 2/4
45	2049	5	1803	4 + 2/4	1557	4
50	2217	5 + 2/4	1951	5	1685	4 + 1/4
55	2382	6	2096	5 + 1/4	1810	4 + 2/4
60	2542	6 + 1/4	2237	5 + 2/4	1932	4 + 3/4
70	2854	7 + 1/4	2511	6 + 1/4	2169	5 + 2/4
80	3154	8	2776	7	2397	6

When a food allergy or a food intolerance is suspected, the dog's diet could be changed without any food transition. The hypoallergenic nutritional management may be continued for the pet's lifetime. It is recommended to divide the daily intake into 2 meals. For the mixed feeding table please see the end of the section "Adverse reaction to food".

NOTE: For the mixed feeding table please see the end of the section Dermatology.

## ADVERSE REACTION TO FOOD



### RECOMMENDED FOR CASES OF

- Adverse Food Reactions (AFR) with dermatologic and/or gastrointestinal signs:
  - diagnosis: food elimination trial
  - management
- Canine Atopic Dermatitis (CAD) associated with AFR
- Chronic diarrhea
- Inflammatory Bowel Disease (IBD)
- Exocrine Pancreatic Insufficiency (EPI)
- Small intestinal bacterial overgrowth (SIBO)

### NOT RECOMMENDED IN CASE OF

- Pancreatitis
- Growth, gestation/lactation



#### HYDROLYSED PROTEIN

Hydrolysed protein with low molecular weight to ensure the food is hypoallergenic.



#### SKIN BARRIER

Formulated to support the skin's natural protective barrier for optimal skin health.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	64.8 g	6.4 g
Fat content	35.4 g	3.5 g
Carbohydrate	127.5 g	12.6 g
Crude fibre	15.2 g	1.5 g
Dietary fibre	22.3 g	2.2 g
Calcium	2 g	0.2 g
Phosphorus	1.6 g	0.16 g
Potassium	2 g	0.2 g
Sodium	2.4 g	0.24 g
Chloride	2 g	0.2 g
Magnesium	0.2 g	0.02 g
Copper	5.1 mg	0.5 mg
Zinc	45.5 mg	4.5 mg
Taurine	1.2 g	0.12 g
Vitamin A	10121.5 IU	1000 IU
Vitamin E	151.18 mg	15 mg
Omega 6	16.2 g	1.6 g
Omega 3	1.7 g	0.17 g
EPA + DHA	1.2 g	0.12 g
Metabolisable energy*		99 kcal/100 g

\*Calculated according to NRC2006.

### COMPOSITION

Derivatives of vegetable origin, vegetable protein extracts, oils and fats, meat and animal derivatives, minerals. Selected sources of protein: hydrolysed soya protein concentrate, hydrolysed poultry by-products. Selected source of carbohydrate: pea starch.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin A: 3000 IU, Vitamin D3: 320 IU, E1 (Iron): 11 mg, E2 (Iodine): 1 mg, E4 (Copper): 3.8 mg, E5 (Manganese): 15 mg, E6 (Zinc): 36 mg - Technological additives: Clinoptilolite of sedimentary origin: 2.5 g.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

## HYPOALLERGENIC



## HYPOALLERGENIC

### MODERATE CALORIE



Available in 1.5 kg, 7 kg, 14 kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION			
		NORMAL		OVERWEIGHT	
		grams	cups	grams	cups
	2	43	4/8	36	3/8
	4	72	7/8	61	6/8
	6	98	1 + 1/8	82	7/8
	8	121	1 + 3/8	102	1 + 1/8
	10	143	1 + 5/8	121	1 + 3/8
	15	194	2 + 2/8	163	1 + 7/8
	20	241	2 + 6/8	203	2 + 3/8
	25	285	3 + 2/8	240	2 + 6/8
	30	326	3 + 6/8	275	3 + 1/8
	35	366	4 + 1/8	308	3 + 4/8
	40	405	4 + 5/8	341	3 + 7/8
	45	442	5	372	4 + 2/8
	50	479	5 + 4/8	403	4 + 5/8
	55	514	5 + 7/8	433	5
	60	549	6 + 2/8	462	5 + 2/8
	70	616	7	519	5 + 7/8
	80	681	7 + 6/8	573	6 + 4/8

When a food allergy or a food intolerance is suspected, the dog's diet could be changed without any food transition. Nutritional management with Hypoallergenic Moderate Calorie may be continued for the pet's lifetime. It is recommended to divide the daily intake into 2 meals.

## ADVERSE REACTION TO FOOD



### RECOMMENDED FOR DOGS WITH TENDENCY TO BE OVERWEIGHT IN CASES OF

- Adverse Food Reactions (AFR) with dermatologic and/or gastrointestinal signs:
  - diagnosis: food elimination trial
  - management
- Canine Atopic Dermatitis (CAD) + chronic pruritus
- Chronic diarrhea
- Inflammatory Bowel Disease (IBD)
- Exocrine Pancreatic Insufficiency (EPI)
- Small intestinal bacterial overgrowth (SIBO)

### NOT RECOMMENDED IN CASE OF

- Acute pancreatitis\*
- Gestation/lactation/growth

\*may be fed in case of chronic pancreatitis



#### HYDROLYSED PROTEIN

Hydrolysed protein with low molecular weight to ensure the food is hypoallergenic.



#### MODERATE CALORIE

Moderate calorie content to help maintain ideal weight.



#### SKIN BARRIER

Formulated to support the skin's natural protective barrier for optimal skin health.



#### MODERATE PHOSPHORUS

A moderate phosphorus intake to help support healthy kidney function.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	61.6 g	23 g
Fat content	32.1 g	12 g
Carbohydrate	127.8 g	47.7 g
Crude fibre	2.9 g	1.1 g
Dietary fibre	13.9 g	5.2 g
Calcium	2.2 g	0.81 g
Phosphorus	1.6 g	0.6 g
Potassium	2.1 g	0.8 g
Sodium	1.1 g	0.4 g
Chloride	1.4 g	0.53 g
Magnesium	0.2 g	0.07 g
Copper	4 mg	1.5 mg
Zinc	45.8 mg	17.1 mg
Taurine	0.6 g	0.21 g
Vitamin A	7902.5 IU	2950 IU
Vitamin E	160.7 mg	60 mg
Omega 6	8.7 g	3.25 g
Omega 3	2.3 g	0.87 g
EPA + DHA	1 g	0.37 g
Metabolisable energy*		373 kcal/100 g

\*Calculated according to NRC2006.

### COMPOSITION

Rice, hydrolysed soya protein isolate, animal fats, hydrolysed poultry liver, minerals, beet pulp, soya oil, fish oil, fructo-oligosaccharides, borage oil, marigold extract (source of lutein). Protein sources: hydrolysed soya protein isolate, hydrolysed poultry liver. Carbohydrate source: rice.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin A: 27000 IU, Vitamin D3: 800 IU, E1 (Iron): 41 mg, E2 (Iodine): 3.7 mg, E4 (Copper): 15 mg, E5 (Manganese): 54 mg, E6 (Zinc): 153 mg, E8 (Selenium): 0.26 mg - Technological additives: Clinoptilolite of sedimentary origin: 10 g - Preservatives - Antioxidants.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



## HYPOALLERGENIC

SMALL DOGS



Available in 1 kg, 3.5 kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	cups	grams	cups	grams	cups
	2	54	5/8	48	1/2	41	3/8
	2.5	64	5/8	56	5/8	49	4/8
	3	73	6/8	64	5/8	56	5/8
	3.5	82	7/8	72	6/8	62	5/8
	4	91	1	80	7/8	69	6/8
	5	107	1 + 1/8	94	1	82	7/8
	6	123	1 + 2/8	108	1 + 1/8	94	1
	7	138	1 + 4/8	122	1 + 2/8	105	1 + 1/8
	8	153	1 + 5/8	134	1 + 3/8	116	1 + 2/8
	9	167	1 + 6/8	147	1 + 4/8	127	1 + 3/8
	10	181	1 + 7/8	159	1 + 5/8	137	1 + 3/8

When a food allergy or a food intolerance is suspected, the dog's diet could be changed without any food transition. Nutritional management with Hypoallergenic Small Dog may be continued for the pet's lifetime. It is recommended to divide the daily intake into 2 meals.

NOTE: For the mixed feeding table please see the end of the section Dermatology.

## ADVERSE REACTION TO FOOD



### RECOMMENDED FOR ADULT SMALL BREED DOGS (< 10KG) IN CASE OF

- Adverse Food Reactions (AFR) with dermatologic and/or gastro-intestinal signs:
  - diagnosis: food elimination trial
  - management
- Canine Atopic Dermatitis (CAD) + chronic pruritus
- Chronic diarrhea
- Inflammatory Bowel Disease (IBD)
- Exocrine Pancreatic Insufficiency (EPI)
- Small intestinal bacterial overgrowth (SIBO)

### NOT RECOMMENDED IN CASE OF

- Pancreatitis
- Gestation/lactation/growth



#### HYDROLYSED PROTEIN

Hydrolysed protein with low molecular weight to ensure the food is hypoallergenic.



#### LOW RSS

Helps lower the concentration of ions contributing to crystal formation.



#### SKIN BARRIER

Formulated to support the skin's natural protective barrier for optimal skin health.



#### DENTAL HEALTH

Contains specific nutrients which contribute to maintaining good oral health.

[S/O®] This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	61.6 g	24 g
Fat content	41.1 g	16 g
Carbohydrate	106.3 g	41.4 g
Crude fibre	1.8 g	0.7 g
Dietary fibre	11.3 g	4.4 g
Calcium	2.1 g	0.8 g
Phosphorus	1.6 g	0.61 g
Potassium	2.3 g	0.9 g
Sodium	2.6 g	1 g
Chloride	3.8 g	1.47 g
Magnesium	0.2 g	0.07 g
Copper	3.9 mg	1.5 mg
Zinc	43.9 mg	17.1 mg
Taurine	0.5 g	0.21 g
Vitamin A	7577.7 IU	2950 IU
Vitamin E	154.1 mg	60 mg
Omega 6	10 g	3.91 g
Omega 3	2.4 g	0.92 g
EPA + DHA	0.9 g	0.35 g
Metabolisable energy*		389 kcal/100 g

\*Calculated according to NRC2006.

### COMPOSITION

Rice, hydrolysed soya protein isolate, animal fats, minerals, hydrolysed poultry liver, soya oil, beet pulp, fish oil, fructo-oligosaccharides, borage oil, marigold extract (source of lutein). Protein sources: hydrolysed soya protein isolate, hydrolysed poultry liver. Carbohydrate source: rice.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin A: 27000 IU, Vitamin D3: 800 IU, E1 (Iron): 41 mg, E2 (Iodine): 3.7 mg, E4 (Copper): 15 mg, E5 (Manganese): 54 mg, E6 (Zinc): 153 mg, E8 (Selenium): 0.26 mg - Technological additives: Clinoptilolite of sedimentary origin: 10 g - Preservatives - Antioxidants.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



HYPOALLERGENIC MODERATE CALORIE



S

SMALL DOGS

HYPOALLERGENIC

HEALTH MANAGEMENT VITAL SUPPORT GASTROINTESTINAL TRACT WEIGHT MANAGEMENT



## SENSITIVITY CONTROL



Available in 1.5 kg, 7 kg, 14 kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

DOG'S WEIGHT (kg)	BODY CONDITION					
	THIN		NORMAL		OVERWEIGHT	
	grams	cups	grams	cups	grams	cups
2	64	6/8	56	5/8	48	4/8
4	107	1 + 2/8	94	1	81	7/8
6	145	1 + 5/8	127	1 + 3/8	110	1 + 2/8
8	180	2	158	1 + 6/8	137	1 + 4/8
10	212	2 + 3/8	187	2 + 1/8	161	1 + 7/8
15	288	3 + 2/8	253	2 + 7/8	219	2 + 4/8
20	357	4	314	3 + 4/8	271	3
25	422	4 + 6/8	372	4 + 2/8	321	3 + 5/8
30	484	5 + 4/8	426	4 + 6/8	368	4 + 1/8
35	543	6 + 1/8	478	5 + 3/8	413	4 + 5/8
40	601	6 + 6/8	529	6	456	5 + 1/8
45	656	7 + 3/8	577	6 + 4/8	499	5 + 5/8
50	710	8	625	7	540	6 + 1/8
55	763	8 + 5/8	671	7 + 4/8	580	6 + 4/8
60	814	9 + 1/8	716	8 + 1/8	619	7
70	914	10 + 2/8	804	9	695	7 + 7/8
80	1010	11 + 3/8	889	10	768	8 + 5/8

When a food allergy or a food intolerance is suspected, the dog's diet could be changed without any food transition. The nutritional management may be continued for the pet's lifetime. It is recommended to divide the daily intake into 2 meals.

## ADVERSE REACTION TO FOOD



### RECOMMENDED FOR CASES OF

- Adverse Food Reactions (AFR) with dermatologic and/or gastrointestinal signs:
  - diagnosis: food elimination trial
  - management
- Canine Atopic Dermatitis (CAD) + chronic pruritus
- Chronic diarrhea

### NOT RECOMMENDED IN CASE OF

- Acute pancreatitis\*
- Gestation/lactation/growth

\*may be fed in case of chronic pancreatitis



### SELECTED PROTEIN

A limited number of protein sources helps reduce the risk of food allergies.



### SKIN BARRIER

Formulated to support the skin's natural protective barrier for optimal skin health.



### EPA/DHA

Fatty acids to help maintain a healthy digestive system and a healthy skin.



### DIGESTIVE SECURITY

Nutrients which support a balanced digestive system.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	63.4 g	21 g
Fat content	27.2 g	9 g
Carbohydrate	145 g	48 g
Crude fibre	13.6 g	4.5 g
Dietary fibre	29.6 g	9.8 g
Calcium	4.2 g	1.38 g
Phosphorus	2.7 g	0.91 g
Potassium	2.7 g	0.9 g
Sodium	1.2 g	0.4 g
Chloride	3.3 g	1.1 g
Magnesium	0.2 g	0.08 g
Copper	4.5 mg	1.5 mg
Zinc	55.3 mg	18.3 mg
Taurine	0.7 g	0.23 g
Vitamin A	8610.3 IU	2850 IU
Vitamin E	178.2 mg	59 mg
Omega 6	5.6 g	1.87 g
Omega 3	2.3 g	0.75 g
EPA + DHA	1.1 g	0.35 g
Metabolisable energy**		331 kcal/100 g

\*\*Calculated according to NRC2006.

### COMPOSITION

Tapioca, dehydrated duck protein, vegetable fibres, hydrolysed poultry proteins, animal fats, beet pulp, fish oil, minerals, soya oil, psyllium husks and seeds, fructo-oligo-saccharides, marigold extract (source of lutein). Protein sources: dehydrated duck protein, hydrolysed poultry proteins. Carbohydrate source: tapioca.

### ADDITIVES (KG)\*\*\*

Nutritional additives: Vitamin A: 26500 IU, Vitamin D3: 800 IU, E1 (Iron): 40 mg, E2 (Iodine): 3.6 mg, E4 (Copper): 15 mg, E5 (Manganese): 53 mg, E6 (Zinc): 150 mg, E8 (Selenium): 0.25 mg - Preservatives - Antioxidants.

\*\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

## SENSITIVITY CONTROL

### CHICKEN WITH RICE



LOAF

Available in 420 g can

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

DOG'S WEIGHT (kg)	BODY CONDITION					
	THIN		NORMAL		OVERWEIGHT	
	grams	can	grams	can	grams	can
2	161	1/2	142	1/4	123	1/4
4	271	3/4	239	2/4	206	2/4
6	367	3/4	323	3/4	279	3/4
8	456	1	401	1	347	3/4
10	539	1 + 1/4	474	1 + 1/4	410	1
15	731	1 + 3/4	643	1 + 2/4	555	1 + 1/4
20	907	2 + 1/4	798	2	689	1 + 3/4
25	1072	2 + 2/4	943	2 + 1/4	815	2
30	1229	3	1081	2 + 2/4	934	2 + 1/4
35	1379	3 + 1/4	1214	3	1048	2 + 2/4
40	1525	3 + 3/4	1342	3 + 1/4	1159	2 + 3/4
45	1665	4	1466	3 + 2/4	1266	3
50	1802	4 + 1/4	1586	3 + 3/4	1370	3 + 1/4
55	1936	4 + 2/4	1704	4	1471	3 + 2/4
60	2067	5	1819	4 + 1/4	1571	3 + 3/4
70	2320	5 + 2/4	2041	4 + 3/4	1763	4 + 1/4
80	2564	6	2256	5 + 1/4	1949	4 + 3/4

When a food allergy or intolerance is suspected, the dog's diet could be changed without any food transition period. Nutritional management with Sensitivity Control Canine may be continued for the pet's lifetime. It is recommended to divide the daily intake into 2 meals.

NOTE: For the mixed feeding table please see the end of the section Dermatology.

## ADVERSE REACTION TO FOOD



### RECOMMENDED FOR CASES OF

- Adverse Food Reactions (AFR) with dermatologic and/or gastro-intestinal signs:
  - diagnosis: food elimination trial
  - management
- Canine Atopic Dermatitis (CAD) + chronic pruritus
- Chronic diarrhea

### NOT RECOMMENDED IN CASE OF

- Acute pancreatitis\*

\*may be fed in case of chronic pancreatitis



### SELECTED PROTEIN

A limited number of protein sources helps reduce the risk of food allergies.



### SKIN BARRIER

Formulated to support the skin's natural protective barrier for optimal skin health.



### EPA/DHA

Fatty acids to help maintain a healthy digestive system and a healthy skin.



### DIGESTIVE HEALTH

Adapted nutrients support a balanced digestive system.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	73.4 g	9.5 g
Fat content	63.1 g	6 g
Carbohydrate	50.7 g	9.6 g
Crude fibre	9.3 g	1.5 g
Dietary fibre	14.5 g	2.2 g
Calcium	2.1 g	0.41 g
Phosphorus	2 g	0.33 g
Potassium	2 g	0.2 g
Sodium	1.6 g	0.16 g
Chloride	2.1 g	0.16 g
Magnesium	0.2 g	0.03 g
Copper	4.1 mg	0.41 mg
Zinc	32.1 mg	3.7 mg
Taurine	1.3 g	0.15 g
Vitamin A	36194.4 IU	2450 IU
Vitamin E	124.1 mg	27 mg
Omega 6	19.6 g	1.8 g
Omega 3	2.4 g	0.27 g
EPA + DHA	1.2 g	0.16 g
Metabolisable energy**		121 kcal/100 g

\*\*Calculated according to NRC2006.

### COMPOSITION

Meat and animal derivatives, cereals, derivatives of vegetable origin, oils and fats, minerals, various sugars. Selected source of protein: chicken (55%). Selected source of carbohydrate: rice (8%).

### ADDITIVES (KG)\*\*\*

Nutritional additives: Vitamin D3: 270 IU, E1 (Iron): 6 mg, E2 (Iodine): 0.4 mg, E4 (Copper): 3.5 mg, E5 (Manganese): 1.7 mg, E6 (Zinc): 17 mg - Technological additives: Clinoptilolite of sedimentary origin: 2.5 g.

\*\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

## SENSITIVITY CONTROL

### CHICKEN WITH RICE

## SENSITIVITY CONTROL

## SENSITIVITY CONTROL

DUCK WITH RICE



LOAF



Available in 420 g can

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	can	grams	can	grams	can
	2	141	1/4	124	1/4	107	1/4
	4	237	1/2	209	1/2	180	1/2
	6	321	3/4	283	3/4	244	1/2
	8	399	1	351	3/4	303	3/4
	10	471	1	415	1	358	3/4
	15	639	1 + 2/4	562	1 + 1/4	485	1 + 1/4
	20	792	2	697	1 + 3/4	602	1 + 2/4
	25	937	2 + 1/4	824	2	712	1 + 3/4
	30	1074	2 + 2/4	945	2 + 1/4	816	2
	35	1206	2 + 3/4	1061	2 + 2/4	916	2 + 1/4
	40	1333	3 + 1/4	1173	2 + 3/4	1013	2 + 2/4
	45	1456	3 + 2/4	1281	3	1106	2 + 3/4
	50	1575	3 + 3/4	1386	3 + 1/4	1197	2 + 3/4
	55	1692	4	1489	3 + 2/4	1286	3
	60	1806	4 + 1/4	1589	3 + 3/4	1373	3 + 1/4
	70	2028	4 + 3/4	1784	4 + 1/4	1541	3 + 3/4
	80	2241	5 + 1/4	1972	4 + 3/4	1703	4

When a food allergy or intolerance is suspected, the dog's diet could be changed without any food transition period. Nutritional management with Sensitivity Control Canine may be continued for the pet's lifetime. It is recommended to divide the daily intake into 2 meals.

## ADVERSE REACTION TO FOOD



### RECOMMENDED FOR CASES OF

- Adverse Food Reactions (AFR) with dermatologic and/or gastro-intestinal signs:
  - diagnosis: food elimination trial
  - management
- Canine Atopic Dermatitis (CAD) + chronic pruritus
- Chronic diarrhea

### NOT RECOMMENDED IN CASE OF

- Acute pancreatitis\*
  - Growth, gestation/lactation
- \*may be fed in case of chronic pancreatitis



#### SELECTED PROTEIN

A limited number of protein sources helps reduce the risk of food allergies.



#### SKIN BARRIER

Formulated to support the skin's natural protective barrier for optimal skin health.



#### EPA/DHA

Fatty acids to help maintain a healthy digestive system and a healthy skin.



#### DIGESTIVE HEALTH

Adapted nutrients support a balanced digestive system.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	65.1 g	9 g
Fat content	50.6 g	7 g
Carbohydrate	88.2 g	12.2 g
Crude fibre	8.7 g	1.2 g
Dietary fibre	10.8 g	1.5 g
Calcium	2.5 g	0.35 g
Phosphorus	2.2 g	0.31 g
Potassium	1.7 g	0.33 g
Sodium	1.1 g	0.15 g
Chloride	0.8 g	0.11 g
Magnesium	0.2 g	0.03 g
Copper	4.3 mg	0.6 mg
Zinc	39.8 mg	5.5 mg
Taurine	1.2 g	0.17 g
Vitamin A	10846 IU	1500 IU
Vitamin E	122.9 mg	17 mg
Omega 6	15.9 g	2.2 g
Omega 3	1.4 g	0.2 g
EPA + DHA	1.2 g	0.17 g
Metabolisable energy**		138 kcal/100 g

\*\*Calculated according to NRC2006.

### COMPOSITION

Meat and animal derivatives, cereals, oils and fats, derivatives of vegetable origin, minerals, various sugars. Selected source of protein: duck (51%). Selected source of carbohydrate: rice (15%).

### ADDITIVES (KG)\*\*\*

Nutritional additives: Vitamin A: 2500 IU, Vitamin D3: 220 IU, E1 (Iron): 8 mg, E2 (Iodine): 0.7 mg, E4 (Copper): 2.5 mg, E5 (Manganese): 10 mg, E6 (Zinc): 38 mg - Technological additives: Clinoptilolite of sedimentary origin: 2.5 g.

\*\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

## SKIN CARE



Available in 2 kg, 8 kg, 11 kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	cups	grams	cups	grams	cups
	2	54	4/8	47	4/8	41	3/8
	4	91	7/8	80	3/4	69	5/8
	6	123	1 + 1/8	108	1	93	7/8
	8	152	1 + 4/8	134	1 + 1/4	116	1 + 1/8
	10	180	1 + 6/8	159	1 + 4/8	137	1 + 2/8
	15	244	2 + 2/8	215	2	186	1 + 6/8
	20	303	2 + 7/8	267	2 + 4/8	230	2 + 1/8
	25	358	3 + 3/8	315	3	272	2 + 5/8
	30	411	3 + 7/8	362	3 + 3/8	312	3
	35	461	4 + 3/8	406	3 + 7/8	351	3 + 3/8
	40	510	4 + 7/8	449	4 + 2/8	387	3 + 5/8
	45	557	5 + 2/8	490	4 + 5/8	423	4
	50	603	5 + 6/8	530	5	458	4 + 3/8
	55	570	5 + 3/8	647	6 + 1/8	492	4 + 5/8
	60	691	6 + 4/8	608	5 + 6/8	525	5
	70	776	7 + 3/8	683	6 + 4/8	589	5 + 5/8
	80	857	8 + 1/8	754	7 + 1/8	652	6 + 1/8

As soon as the first signs of skin disease appear, the diet should be prescribed for a 2-month period to obtain visible results. For chronic disease, the dermatological diet could be maintained for the dog's entire lifetime. It is recommended to divide the daily intake into 2 meals.

## DERMATOSIS & HAIR LOSS



### RECOMMENDED FOR CASES OF

- Canine atopic dermatitis (skin lesions and otitis externa)
- Skin and hair support in case of other dermatitis\* such as:
  - Flea Bite Allergic dermatitis
  - Pyodermatitis

### NOT RECOMMENDED IN CASE OF

- Growth, gestation/lactation
- \* in association with antiparasitic/antibiotic treatment



#### DERMAL CARE

Helps in the nutritional management of dogs with reactive skin.



#### SKIN BARRIER

Formulated to support the skin's natural protective barrier for optimal skin health.



#### DIGESTIVE TOLERANCE

Nutrients which support a balanced intestinal flora and digestive transit.



#### ANTIOXIDANT COMPLEX

A synergistic complex of antioxidants to help neutralise free radicals.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	59 g	23 g
Fat content	41 g	16 g
Carbohydrate	112.8 g	44 g
Crude fibre	5.1 g	2 g
Dietary fibre	19 g	7.4 g
Calcium	1.9 g	0.75 g
Phosphorus	1.4 g	0.55 g
Potassium	1.8 g	0.7 g
Sodium	0.8 g	0.3 g
Chloride	2.7 g	1.04 g
Magnesium	0.2 g	0.08 g
Copper	3.8 mg	1.5 mg
Zinc	43.6 mg	17 mg
Taurine	0.5 g	0.2 g
Vitamin A	7948.7 IU	3100 IU
Vitamin E	153.8 mg	60 mg
Omega 6	8.3 g	3.23 g
Omega 3	2.8 g	1.09 g
EPA + DHA	1 g	0.4 g
Metabolisable energy**		390 kcal/100 g

\*\*Calculated according to NRC2006.

### COMPOSITION

Maize, wheat gluten\*\*\*, rice, tapioca, animal fats, maize gluten, hydrolysed animal proteins, minerals, beet pulp, soya oil, fish oil, vegetable fibres, flax seeds, fructo-oligo-saccharides, mono and diglycerides of palmitic and stearic acids esterified with citric acid, borage oil, marigold extract (source of lutein).

\*\*\*L.I.P.: protein selected for its very high digestibility

### ADDITIVES (KG)\*\*\*

Nutritional additives: Vitamin A: 30000 IU, Vitamin D3: 800 IU, E1 (Iron): 74 mg, E2 (Iodine): 5.9 mg, E4 (Copper): 14 mg, E5 (Manganese): 71 mg, E6 (Zinc): 152 mg, E8 (Selenium): 0.19 mg - Technological additives: Clinoptilolite of sedimentary origin: 5 g - Preservatives - Antioxidants.

\*\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

## SKIN CARE

DUCK WITH RICE

SENSITIVITY CONTROL



WEIGHT MANAGEMENT

GASTROINTESTINAL TRACT

VITAL SUPPORT

HEALTH MANAGEMENT



## SKIN CARE

SMALL DOGS

S



Available in 2 kg, 4 kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	cups	grams	cups	grams	cups
	2	53	5/8	47	4/8	41	4/8
	2.5	63	5/8	56	5/8	48	4/8
	3	72	6/8	64	5/8	55	5/8
	3.5	81	7/8	71	6/8	62	5/8
	4	90	1	79	7/8	68	6/8
	5	106	1 + 1/8	93	1	81	7/8
	6	122	1 + 2/8	107	1 + 1/8	93	1
	7	137	1 + 4/8	120	1 + 2/8	104	1 + 1/8
	8	151	1 + 5/8	133	1 + 3/8	115	1 + 2/8
	9	165	1 + 6/8	145	1 + 4/8	125	1 + 3/8
	10	179	1 + 7/8	157	1 + 5/8	136	1 + 4/8

As soon as the first signs of skin disease appear, the diet should be prescribed for a 2-month period to obtain visible results. For chronic disease, the dermatological diet can be maintained for the dog's entire lifetime. It is recommended to divide the daily intake into 2 meals.

## DERMATOSIS & HAIR LOSS



S

SMALL DOGS

### RECOMMENDED FOR ADULT SMALL BREED DOGS (ADULT WEIGHT <10KG) IN CASES OF

- Canine atopic dermatitis (skin lesions and otitis externa)
- Skin and hair support in case of other dermatitis\* such as:
  - Flea Bite Allergic dermatitis
  - Pyodermatitis

### NOT RECOMMENDED IN CASE OF

- Growth, gestation/lactation



#### DERMAL CARE

Helps in the nutritional management of dogs with reactive skin.



#### SKIN BARRIER

Formulated to support the skin's natural protective barrier for optimal skin health.



#### ANTIOXIDANT COMPLEX

A synergistic complex of antioxidants to help neutralise free radicals.



#### DENTAL HEALTH

Contains specific nutrients which contribute to maintaining good oral health.

[S/O\*] This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	63.5 g	25 g
Fat content	40.6 g	16 g
Carbohydrate	106.9 g	42.1 g
Crude fibre	3.3 g	1.3 g
Dietary fibre	15.2 g	6 g
Calcium	1.9 g	0.74 g
Phosphorus	1.4 g	0.55 g
Potassium	1.8 g	0.7 g
Sodium	1 g	0.4 g
Chloride	2.6 g	1.03 g
Magnesium	0.2 g	0.09 g
Copper	3.8 mg	1.5 mg
Zinc	43.2 mg	17 mg
Taurine	0.5 g	0.2 g
Vitamin A	7874 IU	3100 IU
Vitamin E	152.4 mg	60 mg
Omega 6	9.2 g	3.64 g
Omega 3	2.9 g	1.14 g
EPA + DHA	1 g	0.4 g
Metabolisable energy**		394 kcal/100 g

\*\*Calculated according to NRC2006.

### COMPOSITION

Maize, wheat gluten\*\*\*, rice, tapioca, animal fats, maize gluten, minerals, hydrolysed animal proteins, chicory pulp, soya oil, fish oil, flax seeds, fructo-oligo-saccharides, mono - and diglycerides of palmitic and stearic acids esterified with citric acid, borage oil, marigold extract (source of lutein).

\*\*\*L.I.P.: protein selected for its very high digestibility

### ADDITIVES (KG)\*\*\*\*

Nutritional additives: Vitamin A: 30000 IU, Vitamin D3: 800 IU, E1 (Iron): 72 mg, E2 (Iodine): 5.7 mg, E4 (Copper): 14 mg, E5 (Manganese): 70 mg, E6 (Zinc): 151 mg, E8 (Selenium): 0.19 mg - Technological additives: Clinoptilolite of sedimentary origin: 5 g - Preservatives - Antioxidants.

\*\*\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



## SKIN CARE PUPPY

SMALL DOGS

S



Available in 2 kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

PUPPY'S AGE (months)	ADULT WEIGHT					
	2 kg		5 kg		10 kg	
	grams	cups	grams	cups	grams	cups
2	50	3/8	95	6/8	148	1 + 2/8
3	55	4/8	107	7/8	174	1 + 4/8
4	56	4/8	111	7/8	185	1 + 4/8
5	56	4/8	112	7/8	188	1 + 5/8
6	49	3/8	101	7/8	187	1 + 4/8
7	41	3/8	91	6/8	170	1 + 3/8
8	40	3/8	81	5/8	152	1 + 2/8
9	40	3/8	80	5/8	135	1 + 1/8
10	40	3/8	79	5/8	134	1 + 1/8

As soon as the first signs of skin disease appear, the diet should be prescribed for a 2-month period to obtain visible results. For chronic disease, the dermatological diet can be maintained for the dog's entire lifetime. It is recommended to divide the daily intake into 2 meals.

## DERMATOSIS & HAIR LOSS



S

SMALL DOGS

### RECOMMENDED FOR JUNIOR SMALL BREED DOGS (ADULT WEIGHT <10KG) IN CASES OF

- Canine atopic dermatitis (skin lesions and otitis externa)
- Skin and hair support in case of other dermatitis\* such as:
  - Flea Bite Allergic dermatitis
  - Pyodermatitis

\*In association with antiparasitic/antibiotic treatment



#### DERMAL CARE

Helps in the nutritional management of dogs with reactive skin.



#### SKIN BARRIER

Formulated to support the skin's natural protective barrier for optimal skin health.



#### DIGESTIVE HEALTH

Nutrients which support a balanced digestive system.



#### NATURAL DEFENCES SUPPORT

A synergistic complex of antioxidants to help support natural defences.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	72.2 g	29 g
Fat content	49.8 g	20 g
Carbohydrate	78.4 g	31.5 g
Crude fibre	5 g	2 g
Dietary fibre	17.2 g	6.9 g
Calcium	3 g	1.2 g
Phosphorus	2.2 g	0.9 g
Potassium	2.5 g	1 g
Sodium	1 g	0.4 g
Chloride	3 g	1.22 g
Magnesium	0.1 g	0.06 g
Copper	3.7 mg	1.5 mg
Zinc	42.3 mg	17 mg
Taurine	1.2 g	0.49 g
Vitamin A	7715.3 IU	3100 IU
Vitamin E	149.3 mg	60 mg
Omega 6	10.4 g	4.17 g
Omega 3	3 g	1.2 g
EPA + DHA	1 g	0.4 g
Metabolisable energy*		402 kcal/100 g

\*Calculated according to NRC2006.

### COMPOSITION

Rice, wheat gluten\*\*, animal fats, maize, maize gluten, minerals, hydrolysed animal proteins, soya protein isolate\*\*, beet pulp, fish oil, vegetable fibres, soya oil, flax seeds, fructo-oligo-saccharides, borage oil, hydrolysed yeast (source of manno-oligo-saccharides), marigold extract (source of lutein).

\*\*L.I.P.: protein selected for its very high digestibility

### ADDITIVES (KG)\*\*\*

Nutritional additives: Vitamin A: 30000 IU, Vitamin D3: 800 IU, E1 (Iron): 74 mg, E2 (Iodine): 5.7 mg, E4 (Copper): 13 mg, E5 (Manganese): 69 mg, E6 (Zinc): 152 mg, E8 (Selenium): 0.2 mg - Technological additives: Clinoptilolite of sedimentary origin: 5 g - Preservatives - Antioxidants.

\*\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



SKIN CARE PUPPY

## SKIN SUPPORT



Available in 2 kg, 7 kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

DOG'S WEIGHT (kg)	BODY CONDITION					
	THIN		NORMAL		OVERWEIGHT	
	grams	cups	grams	cups	grams	cups
2	57	5/8	51	4/8	44	3/8
4	97	1	85	7/8	73	6/8
6	131	1 + 2/8	107	1 + 1/8	93	1
8	163	1 + 5/8	143	1 + 3/8	124	1 + 2/8
10	192	1 + 7/8	169	1 + 5/8	146	1 + 4/8
15	260	2 + 5/8	229	2 + 2/8	198	2
20	323	3 + 2/8	284	2 + 7/8	246	2 + 4/8
25	382	3 + 6/8	336	3 + 3/8	290	2 + 7/8
30	438	4 + 3/8	385	3 + 7/8	333	3 + 2/8
35	492	4 + 7/8	433	4 + 2/8	374	3 + 6/8
40	543	5 + 3/8	478	4 + 6/8	413	4 + 1/8
45	594	5 + 7/8	522	5 + 1/8	451	4 + 4/8
50	642	6 + 3/8	565	5 + 5/8	488	4 + 7/8
55	690	6 + 7/8	607	6	524	5 + 4/8
60	736	7 + 2/8	648	6 + 3/8	560	5 + 4/8
70	827	8 + 2/8	728	7 + 2/8	628	6 + 1/4
80	914	9 + 1/8	804	8	695	6 + 7/8

As soon as the first signs of skin disease appear, the diet should be prescribed for a 2-month period to obtain visible results. For chronic disease, the dermatological diet may be maintained for the dog's entire lifetime.  
PRECAUTION FOR USE: Turmeric, a major component of SKIN SUPPORT, is a natural colouring agent (yellow) used in the human food industry. On direct contact, the food could stain clothes and hands. We recommend using a measuring cup and washing your hands after use.

NOTE: For the mixed feeding table please see the end of the section Dermatology.

## DERMATOSIS & HAIR LOSS



### RECOMMENDED FOR CASES OF

- Canine atopic dermatitis
- Wound healing
- Skin and hair support in case of other dermatitis\* such as:
  - Flea Bite Allergic dermatitis
  - Pyodermitis

### NOT RECOMMENDED IN CASE OF

- Gestation/lactation
- \*in association with antiparasitic/antibiotic treatment



#### SKIN SUPPORT

A comple of nutrients to help support the skin's natural defences and support healing.



#### SKIN BARRIER

Formulated to support the skin's natural protective barrier for optimal skin health.



#### EPA/DHA

Fatty acids to help maintain a healthy digestive system and a healthy skin.



#### ANTIOXYDANT COMPLEX

A synergistic complex of antioxydants to help neutralise free radicals.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	62.9 g	23 g
Fat content	41 g	15 g
Carbohydrate	112.1 g	41 g
Crude fibre	9.3 g	3.4 g
Dietary fibre	19.4 g	7.1 g
Calcium	3.3 g	1.2 g
Phosphorus	2.7 g	0.99 g
Potassium	2.2 g	0.8 g
Sodium	1.1 g	0.4 g
Chloride	2.2 g	0.82 g
Magnesium	0.2 g	0.09 g
Copper	4.1 mg	1.5 mg
Zinc	46.5 mg	17 mg
Taurine	0.5 g	0.18 g
Vitamin A	8472.3 IU	3100 IU
Vitamin E	164 mg	60 mg
Omega 6	9.2 g	3.35 g
Omega 3	3.3 g	1.2 g
EPA + DHA	1.6 g	0.6 g
Metabolisable energy**		366 kcal/100 g

\*\*Calculated according to NRC2006.

### COMPOSITION

Rice, dehydrated poultry protein, hydrolysed animal proteins, animal fats, vegetable fibres, fish oil, minerals, beet pulp, soya oil, aloe vera extract (0.06%), borage oil, marigold extract (source of lutein).

### ADDITIVES (KG)\*\*\*

Nutritional additives: Vitamin A: 29500 IU, Vitamin D3: 800 IU, Vitamin C: 280 mg, E1 (Iron): 39 mg, E2 (Iodine): 3.9 mg, E4 (Copper): 12 mg, E5 (Manganese): 51 mg, E6 (Zinc): 129 mg, E8 (Selenium): 0.08 mg - Curcumine extract: 4.4 g - Technological additives: Clinoptilolite of sedimentary origin: 10 g - Preservatives - Antioxidants.

\*\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

## SKIN SUPPORT

# DOGS MIX FEEDING GUIDELINES



### MIX: HYPOALLERGENIC DRY + HYPOALLERGENIC 200G WET

DOG'S WEIGHT (kg)	CAN	WEIGHT CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)
2	1/2+	26	2/8	19	2/8	13	1/8
4	1/2+	61	6/8	50	5/8	40	4/8
6	1+	66	6/8	51	5/8	37	3/8
8	1+	94	1 + 1/8	76	7/8	59	5/8
10	1+	120	1 + 3/8	100	1 + 1/8	79	7/8
15	1+	182	2 + 1/8	154	1 + 6/8	126	1 + 4/8
20	1+	238	2 + 6/8	203	2 + 3/8	168	2
25	1+	291	3 + 3/8	250	2 + 7/8	208	2 + 3/8
30	1+	341	4	294	3 + 3/8	247	2 + 7/8
40	1+	435	5	377	4 + 3/8	318	3 + 5/8
50	1+	524	6 + 1/8	455	5 + 2/8	386	4 + 4/8
60	1+	609	7	529	6 + 1/8	450	5 + 2/8
70	1+	690	8	601	7	512	5 + 7/8
80	1+	768	8 + 7/8	669	7 + 6/8	571	6 + 5/8



### MIX: HYPOALLERGENIC DRY + HYPOALLERGENIC 400G WET

DOG'S WEIGHT (kg)	CAN	WEIGHT CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)
2	1/4+	26	2/8	19	2/8	13	1/8
4	1/4+	61	6/8	50	5/8	40	4/8
6	1/2+	66	6/8	51	5/8	37	3/8
8	1/2+	94	1 + 1/8	76	7/8	59	5/8
10	1/2+	120	1 + 3/8	100	1 + 1/8	79	7/8
15	1+	130	1 + 4/8	102	1 + 1/8	74	7/8
20	1+	186	2 + 1/8	151	1 + 6/8	116	1 + 3/8
25	1+	239	2 + 6/8	198	2 + 2/8	156	1 + 6/8
30	1+	289	3 + 3/8	242	2 + 6/8	195	2 + 2/8
40	1+	383	4 + 4/8	325	3 + 6/8	266	3 + 1/8
50	1+	472	5 + 4/8	403	4 + 5/8	334	3 + 7/8
60	1+	557	6 + 4/8	477	5 + 4/8	398	4 + 5/8
70	1+	638	7 + 3/8	549	6 + 3/8	460	5 + 3/8
80	1+	716	8 + 2/8	617	7 + 1/8	519	6



# DOGS MIX FEEDING GUIDELINES



24H

MIX: SKIN SUPPORT DRY + SENSITIVITY CONTROL  
CHICKEN WITH RICE 420 G WET

DOG'S WEIGHT (kg)	CAN	WEIGHT CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)
2	1/4+	20	2/8	13	1/8	6	0
4	1/4+	59	5/8	48	4/8	36	3/8
6	1/4+	94	7/8	78	6/8	62	5/8
8	1/2+	88	7/8	68	5/8	49	4/8
10	1/2+	117	1 + 1/8	94	7/8	71	6/8
15	1/2+	186	1 + 7/8	154	1 + 4/8	123	1 + 2/8
20	1+	173	1 + 6/8	135	1 + 3/8	96	1
25	1+	232	2 + 2/8	186	1 + 7/8	141	1 + 3/8
30	1+	288	2 + 7/8	236	2 + 3/8	183	1 + 7/8
40	1+	394	3 + 7/8	328	3 + 2/8	263	2 + 5/8
50	1+	493	4 + 7/8	416	4 + 1/8	339	3 + 3/8
60	1+	587	5 + 7/8	498	5	410	4 + 1/8
70	1+	677	6 + 6/8	578	5 + 6/8	479	4 + 6/8
80	1+	764	7 + 5/8	654	6 + 4/8	545	5 + 3/8

# DOGS GROWING FEEDING GUIDELINES



24H

HYPOALLERGENIC DOG

PUPPY'S AGE in months	ADULT WEIGHT					
	2 kg		5 kg		10 kg	
	DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)
6	48	4/8	100	1 + 1/8	185	2 + 1/8
7	40	4/8	90	1	167	1 + 7/8
8	39	4/8	79	7/8	150	1 + 6/8
9	39	4/8	79	7/8	133	1 + 4/8
10	39	4/8	78	7/8	132	1 + 4/8
11	-	-	-	-	131	1 + 4/8
12	-	-	-	-	-	-

	20 kg		30 kg		40 kg	
	DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)
6	312	3 + 5/8	422	4 + 7/8	523	6
7	290	3 + 3/8	423	4 + 7/8	524	6 + 1/8
8	270	3 + 1/8	393	4 + 4/8	492	5 + 6/8
9	247	2 + 7/8	366	4 + 2/8	461	5 + 3/8
10	224	2 + 5/8	334	3 + 7/8	429	5
11	222	2 + 5/8	303	3 + 4/8	403	4 + 5/8
12	220	2 + 4/8	301	3 + 4/8	377	4 + 3/8



# DOGS GROWING FEEDING GUIDELINES



**24H SKIN SUPPORT DOG**

PUPPY'S AGE in months	ADULT WEIGHT					
	2 kg		5 kg		10 kg	
	DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)
6	53	5/8	111	1 + 3/8	206	2 + 4/8
7	45	4/8	100	1 + 2/8	186	2 + 2/8
8	44	4/8	88	1	167	2
9	44	4/8	88	1	149	1 + 6/8
10	44	4/8	87	1	148	1 + 6/8
11	-	-	-	-	146	1 + 6/8
12	-	-	-	-	-	-

	20 kg		30 kg		40 kg	
	DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)
6	348	4 + 1/8	470	5 + 5/8	583	7
7	323	3 + 7/8	471	5 + 5/8	584	7
8	301	3 + 5/8	438	5 + 2/8	548	6 + 4/8
9	275	3 + 2/8	408	4 + 7/8	514	6 + 1/8
10	250	3	372	4 + 3/8	479	5 + 6/8
11	247	3	338	4	449	5 + 3/8
12	246	2 + 7/8	335	4	421	5



**24H MIX: SKIN SUPPORT DOG DRY + SENSITIVITY CONTROL CHICKEN WITH RICE 420G WET**

PUPPY'S AGE in months	ADULT WEIGHT					
	2 kg	5 kg	10 kg	20 kg	30 kg	40 kg
	1/8 CAN	1/4 CAN	1/4 CAN	1/4 CAN	1/4 CAN	1/2 CAN
6	35	74	169	75	115	115
7	26	63	149	68	115	115
8	25	51	130	60	105	103
9	25	51	112	52	95	92
10	25	50	111	44	83	81
11	-	-	10	43	72	71
12	-	-	-	42	71	62





## SATIETY WEIGHT MANAGEMENT



Available in 1.5 kg, 6 kg, 12 kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

IDEAL WEIGHT (kg)*	STAGE		
	START	4 WEEKS AFTER**	MAINTENANCE AFTER WEIGHT LOSS
	grams	grams	grams
10	148	126	139
12	169	145	159
14	190	163	179
16	210	180	198
18	229	197	216
20	248	213	234
25	293	251	277
30	336	288	317
35	378	324	356
40	417	358	393
45	456	391	430
50	493	423	465
55	530	454	500
60	566	485	533
70	635	544	599
80	702	602	662
90	767	657	723

\*For dogs less than 10kg it is recommended to use Satiety Small Dog  
\*\*If the dog did not lose weight

**Individual target weight is determined** and then feeding amount established. The recommended daily intake is based on the pet's ideal weight. The starting ration may need to be adjusted during follow-up visits, based on each individual's response and the rate of weight loss. An objective of losing 1-3% bodyweight per week is generally achievable over the first 3 months, and of 0.5-2% afterwards.

**How is the energy allocation for maintenance after weight loss calculated?** Energy requirements to maintain stable body weight after weight loss are lower than those of dogs that have never been obese. Based on current scientific insights (German & al. 2011), our maintenance after weight loss rations are calculated by increasing the weight loss ration by +10%.

**NOTE: maintenance rations may need to be adjusted based on individual evolution.** German AJ & al. Low-maintenance energy requirements of obese dogs after weight loss. British Journal of Nutrition (2011), 106, S93-S96

## WEIGHT MANAGEMENT



### RECOMMENDED FOR WEIGHT LOSS AND MAINTENANCE AFTER WEIGHT LOSS IN CASE OF

- Overweight/Obesity
- Stabilized Diabetes Mellitus with overweight/obesity
- Hyperlipidemia with overweight/obesity

### NOT RECOMMENDED IN CASE OF

- Dogs with a history of constipation in response to a high fibre diet
- Growth, gestation/lactation



#### EFFECTIVE WEIGHT MANAGEMENT

Provides safe weight loss and helps avoid weight regain. 97% of dogs lost weight in 3 months.



#### BEGGING CONTROL

High natural fiber level keeps dogs satisfied between meals. Helps control begging in 83% of dogs during weight loss.



#### MUSCLE MASS MAINTENANCE

High protein content helps support healthy weight loss while maintaining muscle mass.



### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	112.44 g	30.0 g
Fat content	35.61 g	9.5 g
Digestible Carbohydrates (Starch)	63.34 g	16.9 g
NFE	105.70 g	28.2 g
Crude fibre	63.72 g	17.0 g
Dietary fibre	106.07 g	28.3 g
Omega 6	7.61 g	2.03 g
Omega 3	2.92 g	0.78 g
EPA + DHA	1.50 g	0.40 g
Calcium	3.37 g	0.90 g
Phosphorus	2.62 g	0.70 g
Sodium	1.12 g	0.30 g
Metabolisable energy*		267 kcal/100 g

\*Calculated according to NRC2006TDF.

### COMPOSITION

Vegetable fibres, dehydrated poultry protein, wheat gluten\*, tapioca, maize gluten, hydrolysed animal proteins, wheat, maize, animal fats, beet pulp, fish oil, minerals, psyllium husks and seeds, soya oil, fructo-oligo-saccharides, glucosamine, marigold extract (source of lutein), hydrolysed cartilage (source of chondroitin).

\*L.I.P. (Low Indigestible Protein): protein selected because of its very high assimilation.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin A: 19000 IU, Vitamin D3: 1000 IU, Iron (3b103): 35 mg, Iodine (3b201, 3b202): 3.5 mg, Copper (3b405, 3b406): 11 mg, Manganese (3b502, 3b504): 46 mg, Zinc (3b603, 3b605, 3b606): 137 mg, Selenium (3b801, 3b811, 3b812): 0.06 mg - Preservatives - Antioxidants.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

## SATIETY WEIGHT MANAGEMENT



## SATIETY WEIGHT MANAGEMENT



Available in 195 g, 410 g can

### RECOMMENDED DAILY INTAKE & FEEDING DURATION BASED ON 410G

IDEAL WEIGHT (kg)	STAGE					
	START		4 WEEKS AFTER*		MAINTENANCE AFTER WEIGHT LOSS	
	grams	can 410g	grams	can 410g	grams	can 410g
2	191	2/4	164	2/4	180	2/4
4	321	3/4	275	3/4	303	3/4
6	435	1	373	1	410	1
8	540	1 + 1/4	463	1 + 1/4	509	1 + 1/4
10	638	1 + 2/4	547	1 + 1/4	602	1 + 2/4
15	865	2	741	1 + 3/4	815	2
20	1 073	2 + 2/4	920	2 + 1/4	1 012	2 + 2/4
25	1 268	3	1 087	2 + 3/4	1 196	3
30	1 454	3 + 2/4	1 247	3	1 371	3 + 1/4
35	1 633	4	1 399	3 + 2/4	1 539	3 + 3/4
40	1 805	4 + 2/4	1 547	3 + 3/4	1 701	4 + 1/4
45	1 971	4 + 3/4	1 690	4	1 859	4 + 2/4
50	2 133	5 + 1/4	1 828	4 + 2/4	2 011	5
55	2 291	5 + 2/4	1 964	4 + 3/4	2 160	5 + 1/4
60	2 446	6	2 096	5	2 306	5 + 2/4

\* If the dog did not lose weight

**Individual target weight is determined** and then feeding amount established. The recommended daily intake is based on the pet's ideal weight. The starting ration may need to be adjusted during follow-up visits, based on each individual's response and the rate of weight loss. An objective of losing 1-3% bodyweight per week is generally achievable over the first 3 months, and of 0.5-2% afterwards.

**How is the energy allocation for maintenance after weight loss calculated?** Energy requirements to maintain stable body weight after weight loss are lower than those of dogs that have never been obese. Based on current scientific insights (German & al. 2011), our maintenance after weight loss rations are calculated by increasing the weight loss ration by +10%.

**NOTE: maintenance rations may need to be adjusted based on individual evolution.** German AJ & al. Low-maintenance energy requirements of obese dogs after weight loss. British Journal of Nutrition (2011), 106, S93-S96

## WEIGHT MANAGEMENT



### RECOMMENDED FOR WEIGHT LOSS / MAINTENANCE AFTER WEIGHT LOSS IN CASE OF

- Overweight/Obesity
- Stabilized Diabetes Mellitus with overweight/obesity
- Hyperlipidemia with overweight/obesity

### NOT RECOMMENDED IN CASE OF

- Growth, gestation/lactation



#### EFFECTIVE WEIGHT MANAGEMENT

Helps provide safe weight loss and avoid weight regain.



#### BEGGING CONTROL

Nutritional concept that helps keep dogs satisfied between meals.



#### MUSCLE MASS MAINTENANCE

High protein content helps maintain muscle mass during weight loss.



[S/O] INDEX This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	129.66 g	8.0 g
Fat content	32.41 g	2.0 g
Digestible Carbohydrates (Starch)	38.90 g	2.4 g
NFE	64.83 g	4.0 g
Crude fibre	32.41 g	2.0 g
Dietary fibre	51.86 g	3.2 g
Omega 6	5.67 g	0.35 g
Omega 3	1.46 g	0.09 g
EPA + DHA	0.97 g	0.06 g
Calcium	2.27 g	0.14 g
Phosphorus	1.94 g	0.12 g
Sodium	1.94 g	0.12 g
Metabolisable energy*		62 kcal/100 g

\*Calculated according to NRC2006TDF.

### COMPOSITION

Meat and animal derivatives, derivatives of vegetable origin, cereals, oils and fats, minerals, molluscs and crustaceans.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 290 IU, Iron (3b103): 6.5 mg, Iodine (3b202): 0.2 mg, Copper (3b405, 3b406): 1.8 mg, Manganese (3b502, 3b503, 3b504): 2 mg, Zinc (3b603, 3b605, 3b606): 20 mg - Technological additives: Clinoptilolite of sedimentary origin: 1 g.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

## SATIETY WEIGHT MANAGEMENT



## SATIETY WEIGHT MANAGEMENT

SMALL DOGS

S



Available in 500 g, 1,5 kg, 3 kg, 8 kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT DOG	IDEAL WEIGHT (kg)	STAGE		
		START	4 WEEKS AFTER*	MAINTENANCE AFTER WEIGHT LOSS
		grams	grams	grams
	2	44	37	41
	2.5	52	44	49
	3	59	51	56
	3.5	66	57	63
	4	73	63	69
	4.5	80	69	76
	5	87	74	82
	5.5	93	80	88
	6	100	85	94
	6.5	106	91	100
	7	112	96	105
	7.5	118	101	111
	8	124	106	116
	8.5	129	111	122
	9	135	116	127
	9.5	140	120	132
	10	146	125	138

\* If the dog did not lose weight

**Individual target weight is determined** and then feeding amount established. The recommended daily intake is based on the pet's ideal weight. The starting ration may need to be adjusted during follow-up visits, based on each individual's response and the rate of weight loss. An objective of losing 1-3% bodyweight per week is generally achievable over the first 3 months, and of 0.5-2% afterwards.

**How is the energy allocation for maintenance after weight loss calculated?** Energy requirements to maintain stable body weight after weight loss are lower than those of dogs that have never been obese. Based on current scientific insights [German & al. 2011], our maintenance after weight loss rations are calculated by increasing the weight loss ration by +10%.

**NOTE: maintenance rations may need to be adjusted based on individual evolution.** German AJ & al. Low-maintenance energy requirements of obese dogs after weight loss. British Journal of Nutrition (2011), 106, 593-596.

## WEIGHT MANAGEMENT



### RECOMMENDED FOR WEIGHT LOSS AND MAINTENANCE AFTER WEIGHT LOSS IN CASE OF

- Overweight/Obesity
- Stabilized Diabetes Mellitus with overweight/obesity
- Hyperlipidemia with overweight/obesity

### NOT RECOMMENDED IN CASE OF

- Dogs with a history of constipation in response to a high fibre diet
- Growth, gestation/lactation



#### EFFECTIVE WEIGHT MANAGEMENT

Provides safe weight loss and helps avoid weight regain. 97% of dogs lost weight in 3 months.



#### BEGGING CONTROL

High natural fiber level keeps dogs satisfied between meals. Helps control begging in 83% of dogs during weight loss.



#### SPECIFICALLY FORMULATED FOR SMALL DOGS

Adapted formula to meet the specific digestive, dental and urinary sensitivities of small dogs.

[S/O®]  
INDEX

This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	111.28 g	30.0 g
Fat content	35.24 g	9.5 g
Digestible Carbohydrates (Starch)	61.94 g	16.7 g
NFE	106.08 g	28.6 g
Crude fibre	57.86 g	15.6 g
Dietary fibre	102.00 g	27.5 g
Omega 6	6.82 g	1.84 g
Omega 3	2.30 g	0.62 g
EPA + DHA	1.11 g	0.30 g
Calcium	3.38 g	0.91 g
Phosphorus	2.82 g	0.76 g
Sodium	2.60 g	0.70 g
Metabolisable energy*		270 kcal/100 g

\*Calculated according to NRC2006TDF.

### COMPOSITION

Dehydrated poultry protein, vegetable fibres, wheat gluten\*, tapioca, hydrolysed animal proteins, chicory pulp, wheat, maize, maize gluten, animal fats, minerals, psyllium husks and seeds, fish oil, fatty acid salt, fructo-oligo-saccharides, marigold extract (source of lutein), hydrolysed crustaceans (source of glucosamine), hydrolysed cartilage (source of chondroitin).

\*L.I.P. (Low Indigestible Protein): protein selected because of its very high assimilation.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin A: 19000 IU, Vitamin D3: 1000 IU, Iron (3b103): 34 mg, Iodine (3b201, 3b202): 3.4 mg, Copper (3b405, 3b406): 10 mg, Manganese (3b502, 3b504): 44 mg, Zinc (3b603, 3b605, 3b606): 136 mg, Selenium (3b801, 3b811, 3b812): 0.05 mg - Preservatives - Antioxidants.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



S

SMALL DOGS  
SATIETY WEIGHT MANAGEMENT



## DIABETIC



Available in 1.5 kg, 7 kg, 12 kg bags

### RECOMMENDED DAILY INTAKE & FEEDING DURATION

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	cups	grams	cups	grams	cups
	2	61 g	5/8	54 g	5/8	46 g	4/8
	4	103 g	1 + 1/8	91 g	1	78 g	7/8
	6	139 g	1 + 4/8	123 g	1 + 3/8	106 g	1 + 1/8
	8	173 g	1 + 7/8	152 g	1 + 5/8	131 g	1 + 3/8
	10	205 g	2 + 2/8	180 g	2	155 g	1 + 5/8
	15	277 g	3	244 g	2 + 5/8	211 g	2 + 2/8
	20	344 g	3 + 6/8	303 g	3 + 2/8	261 g	2 + 7/8
	25	407 g	4 + 3/8	358 g	3 + 7/8	309 g	3 + 3/8
	30	466 g	5	410 g	4 + 3/8	354 g	3 + 7/8
	35	523 g	5 + 5/8	461 g	5	398 g	4 + 2/8
	40	578 g	6 + 2/8	509 g	5 + 4/8	440 g	4 + 6/8
	45	632 g	6 + 7/8	556 g	6	480 g	5 + 2/8
	50	684 g	7 + 3/8	602 g	6 + 4/8	520 g	5 + 5/8
	55	735 g	8	646 g	7	558 g	6
	60	784 g	8 + 4/8	690 g	7 + 4/8	596 g	6 + 4/8
	70	880 g	9 + 4/8	775 g	8 + 3/8	669 g	7 + 2/8
	80	973 g	10 + 4/8	856 g	9 + 2/8	739 g	8

DIABETIC is recommended as first line nutritional management for all newly diagnosed cases of Diabetes Mellitus until control of clinical signs and hyperglycemia have been achieved.

**NOTE: The use of this food may affect the dog's sensitivity to insulin. It is highly recommended that blood glucose and insulin response are monitored when starting the diet so that insulin dosage can be adapted.**

## DIABETES MELLITUS




### RECOMMENDED FOR CASES OF

- Diabetes Mellitus and a Body Condition Score (BCS) < 6/9
- Diabetes Mellitus and a BCS > 7/9, until glycemic control has been achieved

### NOT RECOMMENDED IN CASE OF

- Growth, gestation/lactation



#### GLUCOMODULATION

Specific formula to help in the management of post-prandial blood glucose in diabetic dogs.



#### HIGH PROTEIN

High protein content. Maintenance of muscle mass is essential in diabetic dogs.



#### LOW STARCH

Formula that contains a reduced level of starch.

### KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	107.65 g	37.0 g
Fat content	34.91 g	12.0 g
Digestible Carbohydrates (Starch)	55.57 g	19.1 g
NFE	86.41 g	29.7 g
Crude fibre	18.91 g	6.5 g
Dietary fibre	49.75 g	17.1 g
Omega 6	6,20 g	2,13 g
Omega 3	1,86 g	0,64 g
EPA + DHA	0.87 g	0.30 g
Calcium	2,62 g	0,90 g
Phosphorus	2,04 g	0,70 g
Sodium	0.87 g	0.30 g
Metabolisable energy*		344 kcal/100 g

\*Calculated according to NRC2006CF.

### COMPOSITION

Dehydrated poultry protein, barley, wheat gluten\*, maize gluten, vegetable fibres, hydrolysed animal proteins, animal fats, beet pulp, tapioca, fish oil, psyllium husks and seeds, fructo-oligo-saccharides, minerals, marigold extract (source of lutein). Sources of carbohydrates: barley, tapioca.

\*L.I.P. (Low Indigestible Protein): protein selected because of its very high assimilation.

### ADDITIVES (KG)\*\*

Nutritional additives: Vitamin A: 15500 IU, Vitamin D3: 1000 IU, Iron (3b103): 33 mg, Iodine (3b201, 3b202): 3.4 mg, Copper (3b405, 3b406): 10 mg, Manganese (3b502, 3b504): 43 mg, Zinc (3b603, 3b605, 3b606): 125 mg, Selenium (3b801, 3b811, 3b812): 0.05 mg - Preservatives - Antioxidants.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.



DIABETIC

DIABETIC SPECIAL

LOW CARBOHYDRATE



Available in 195 g, 410 g can

RECOMMENDED DAILY INTAKE  
& FEEDING DURATION BASED ON 410g

ADULT DOG	DOG'S WEIGHT (kg)	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		grams	can 410g	grams	can 410g	grams	can 410g
	2	257	3/4	226	2/4	195	2/4
	4	432	1	380	1	328	3/4
	6	585	1 + 2/4	515	1 + 1/4	445	1
	8	726	1 + 3/4	639	1 + 2/4	552	1 + 1/4
	10	858	2	755	1 + 3/4	652	1 + 2/4
	15	1 163	2 + 3/4	1 024	2 + 2/4	884	2 + 1/4
	20	1 443	3 + 2/4	1 270	3	1 097	2 + 3/4
	25	1 706	4 + 1/4	1 502	3 + 3/4	1 297	3 + 1/4
	30	1 956	4 + 3/4	1 722	4 + 1/4	1 487	3 + 3/4
	35	2 196	5 + 1/4	1 933	4 + 3/4	1 669	4
	40	2 428	6	2 136	5 + 1/4	1 845	4 + 2/4
	45	2 652	6 + 2/4	2 334	5 + 3/4	2 015	5
	50	2 870	7	2 525	6 + 1/4	2 181	5 + 1/4
	55	3 082	7 + 2/4	2 713	6 + 2/4	2 343	5 + 3/4
	60	3 290	8	2 895	7	2 501	6
	70	3 694	9	3 250	8	2 807	6 + 3/4
	80	4 083	10	3 593	8 + 3/4	3 103	7 + 2/4

DIABETIC is recommended as first line nutritional management for all newly diagnosed cases of Diabetes Mellitus until control of clinical signs and hyperglycemia have been achieved.

NOTE: The use of this food may affect the dog's sensitivity to insulin. It is highly recommended that blood glucose and insulin response are monitored when starting the diet so that insulin dosage can be adapted.

DIABETES MELLITUS



RECOMMENDED FOR CASES OF

- Diabetes Mellitus and a Body Condition Score (BCS) ≤ 6/9
- Diabetes Mellitus and a BCS > 7/9, until glycemic control has been achieved

NOT RECOMMENDED IN CASE OF

- Growth, gestation/lactation



GLUCOMODULATION

Specific formula to help in the management of post-prandial blood glucose in diabetic dogs.



HIGH PROTEIN

High protein content. Maintenance of muscle mass is essential in diabetic dogs.



LOW STARCH

Formula that contains a reduced level of starch.

[S/O®]  
INDEX]

This product creates an unfavorable environment to creation of oxalate and struvite uroliths in the urine.

KEY VALUES

	Per 1000 kcal	Per 100g as fed
Protein	109.89 g	9.0 g
Fat content	48.84 g	4.0 g
Digestible Carbohydrates (Starch)	19.54 g	1.6 g
NFE	50.06 g	4.1 g
Crude fibre	29.30 g	2.4 g
Dietary fibre	47.62 g	3.9 g
Omega 6	12.21 g	1.00 g
Omega 3	0.73 g	0.06 g
EPA + DHA	0.09 g	0.01 g
Linoleic acid	10,99 g	0,90 g
Calcium	3.42 g	0.28 g
Phosphorus	2.56 g	0.21 g
Sodium	1.22 g	0.10 g
Metabolisable energy*		82 kcal/100 g

\*Calculated according to NRC2006TDF.

COMPOSITION

Meat and animal derivatives, derivatives of vegetable origin, cereals, oils and fats, minerals.

ADDITIVES (KG)\*\*

Nutritional additives: Vitamin D3: 180 IU, Iron (3b103): 3 mg, Iodine (3b201, 3b202): 0.3 mg, Copper (3b405, 3b406): 2 mg, Manganese (3b502, 3b504): 1 mg, Zinc (3b603, 3b605, 3b606): 10 mg.

\*\*Values reflect only levels added to the formula, not those naturally occurring in components of the diet.

DIABETIC





# DOGS MIX FEEDING GUIDELINES



24H

MIX: SATIETY DOG DRY + SATIETY DOG WET CAN

195 G CAN

IDEAL WEIGHT (kg)	CAN	STAGE		
		START	4 WEEKS AFTER*	MAINTENANCE AFTER WEIGHT LOSS
		DRY (g)	DRY (g)	DRY (g)
2	1/2	22	15	19
4	1	29	19	25
6	1	55	41	50
8	1	80	62	73
10	1	102	81	94
15	1	155	126	143
20	1	203	168	189
25	1	248	206	231
30	1	291	243	272
40	1	372	313	348
50	1	448	378	420
60	1	521	440	488

\* If the dog did not lose weight



24H

MIX: SATIETY DOG DRY + SATIETY DOG WET CAN

410 G CAN

IDEAL WEIGHT (kg)	CAN	STAGE		
		START	4 WEEKS AFTER*	MAINTENANCE AFTER WEIGHT LOSS
		DRY (g)	DRY (g)	DRY (g)
2	1/4	20	14	18
4	1/2	27	16	23
6	1/2	53	39	47
8	1/2	77	60	70
10	1	53	32	44
15	1	105	77	94
20	1	153	118	139
25	1	199	157	182
30	1	242	193	222
40	1	322	263	299
50	1	399	328	370
60	1	471	390	438

\* If the dog did not lose weight



24H

MIX: SATIETY SMALL DOG DRY + SATIETY DOG WET CAN

195 G CAN

IDEAL WEIGHT (kg)	CAN	STAGE		
		START	4 WEEKS AFTER*	MAINTENANCE AFTER WEIGHT LOSS
		DRY (g)	DRY (g)	DRY (g)
2	1/2	21	15	19
4	1	29	18	25
6	1	55	41	49
8	1	79	61	72
10	1	101	81	93

\* If the dog did not lose weight



24H

MIX: SATIETY SMALL DOG DRY + SATIETY DOG WET CAN

410 G CAN

IDEAL WEIGHT (kg)	CAN	STAGE		
		START	4 WEEKS AFTER*	MAINTENANCE AFTER WEIGHT LOSS
		DRY (g)	DRY (g)	DRY (g)
2	1/4	20	14	18
4	1/2	27	16	22
6	1/2	53	38	47
8	1/2	77	59	70
10	1	52	31	44


\* If the dog did not lose weight





# DOGS MIX FEEDING GUIDELINES






MIX: DIABETIC DOG DRY +  
DIABETIC LOW CARBOHYDRATE DOG WET CAN

195 G CAN

DOG'S WEIGHT (kg)	CAN	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)
2	1/2	38	3/8	31	3/8	23	2/8
4	1	56	5/8	44	4/8	32	3/8
6	1	93	1	76	7/8	59	5/8
8	1	127	1 + 3/8	106	1 + 1/8	85	7/8
10	1	158	1 + 6/8	134	1 + 4/8	109	1 + 1/8
15	1	231	2 + 4/8	197	2 + 1/8	164	1 + 6/8
20	1	297	3 + 2/8	256	2 + 6/8	215	2 + 3/8
25	1	360	3 + 7/8	311	3 + 3/8	263	2 + 7/8
30	1	420	4 + 4/8	364	3 + 7/8	308	3 + 3/8
40	1	532	5 + 6/8	463	5	393	4 + 2/8
50	1	637	6 + 7/8	555	6	473	5 + 1/8
60	1	738	8	643	7	549	6
70	1	834	9	728	7 + 7/8	622	6 + 6/8
80	1	926	10	810	8 + 6/8	693	7 + 4/8





MIX: DIABETIC DOG DRY +  
DIABETIC LOW CARBOHYDRATE DOG WET CAN

410 G CAN

DOG'S WEIGHT (kg)	CAN	BODY CONDITION					
		THIN		NORMAL		OVERWEIGHT	
		DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)	DRY (g)	CUP (240 cm³)
2	1/4	37	3/8	29	3/8	22	2/8
4	1/2	54	5/8	42	4/8	29	3/8
6	1/2	91	1	74	6/8	57	5/8
8	1/2	124	1 + 3/8	103	1 + 1/8	83	7/8
10	1	107	1 + 1/8	82	7/8	58	5/8
15	1	180	2	146	1 + 5/8	113	1 + 2/8
20	1	246	2 + 5/8	205	2 + 2/8	164	1 + 6/8
25	1	309	3 + 3/8	260	2 + 7/8	211	2 + 2/8
30	1	369	4	313	3 + 3/8	257	2 + 6/8
40	1	481	5 + 2/8	411	4 + 4/8	342	3 + 6/8
50	1	586	6 + 3/8	504	5 + 4/8	422	4 + 5/8
60	1	686	7 + 3/8	592	6 + 3/8	498	5 + 3/8
70	1	782	8 + 4/8	677	7 + 3/8	571	6 + 1/8
80	1	875	9 + 4/8	758	8 + 2/8	642	7



